

# INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C07K 14/705, A61K 38/17, C07K 16/28 A1 (11) International Publication Number:

WO 99/51639

(43) International Publication Date:

14 October 1999 (14.10.99)

(21) International Application Number:

PCT/SE99/00544

(22) International Filing Date:

31 March 1999 (31.03.99)

(30) Priority Data:

9801164-6 9900319-6 2 April 1998 (02.04.98) 28 January 1999 (28.01.99)

SE

(71) Applicant (for all designated States except US): ACTIVE BIOTECH AB [SE/SE]; Scheelevägen 22, S-220 07 Lund (SE).

(72) Inventor; and

LUND-(75) Inventor/Applicant US only): (for GREN-AKERLUND, Evy [SE/SE]; Trollsjövägen 165, S-237 33 Bjärred (SE).

(74) Agent: AWAPATENT AB; P.O. Box 5117, S-200 71 Malmö (SE).

(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published


With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

# (54) Title: AN INTEGRIN HETERODIMER AND A SUBUNIT THEREOF

#### (57) Abstract

A recombinant or isolated integrin heterodimer comprising a novel subunit  $\alpha 10$  in association with a subunit  $\beta$  is described. The  $\alpha$  10 integrin may be purified from bovine chondrocytes on a collagen-type-II affinity column. The integrin or the subunit  $\alpha 10$  can be used as marker or target of all types of cells, e.g. of chondrocytes, osteoblasts and fibroblasts. The integrin or subunit a 10 thereof can be used as marker or target in different physiological or therapeutic methods. They can also be used as active ingredients in pharmaceutical compositions and vaccines.



# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ.	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghans	MG	Madagascar	T.J	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	17M	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG		HU	Hungary	ML	Mali	TT	Trinidad and Tobego
	Bulgaria Benin	IE	Ireland	MIN	Mongolia	UA	Ukraine
BJ	Brazil	IL	Israel	MIR	Mauritania	UG	Uganda
BR		IS	Iceland	MW	Malawi	US	United States of Americ
BY	Belarus	IT	Italy	MX	Mexico	UZ	Uzbekistan
CA	Canada	JP	Japan	NE	Nizer	VN	Viet Nam
CF .	Central African Republic	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CG	Congo	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CH	Switzerland	KP	Democratic People's	NZ	New Zealand		
a	Côte d'Ivoire	KJ	Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic		Liechtenstein	SD	Sudan		
DE	Germany	u		SE	Sweden		
DK	Denmark	LK	Sri Lanka	SG			
EE	Estonia	LR	Liberia	20	Singapore		

WO 99/51639 PCT/SE99/00544

# AN INTEGRIN HETERODIMER AND A SUBUNIT THEREOF

#### FIELD OF THE INVENTION

The present invention relates to a recombinant or isolated integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , the subunit  $\alpha 10$  thereof, homologues and fragments of said integrin and of said subunit  $\alpha 10$  having similar biological activity, processes of producing the same, polynucleotides and oligonucleotides encoding the same, vectors and cells comprising the same, binding entities binding specifically to the same, and the use of the same.

#### BACKGROUND OF THE INVENTION

10

The integrins are a large family of transmembrane glycoproteins that mediate cell-cell and cell-matrix 15 interactions (1-5). All known members of this superfamily are non-covalently associated heterodimers composed of an  $\alpha$ - and a  $\beta$ -subunit. At present, 8  $\beta$ -subunits ( $\beta$ 1- $\beta$ 8) (6) and 16  $\alpha$ -subunits ( $\alpha$ 1- $\alpha$ 9,  $\alpha$ v,  $\alpha$ M,  $\alpha$ L,  $\alpha$ X,  $\alpha$ IIb,  $\alpha$ E and αD) have been characterized (6-21), and these subunits 20 associate to generate more than 20 different integrins. The \$1-subunit has been shown to associate with ten different  $\alpha$ -subunits,  $\alpha 1-\alpha 9$  and  $\alpha v$ , and to mediate interactions with extracellular matrix proteins such as collagens, laminins and fibronectin. The major collagen bind-25 ing integrins are  $\alpha1\beta1$  and  $\alpha2\beta1$  (22-25). The integrins  $\alpha 3\beta 1$  and  $\alpha 9\beta 1$  have also been reported to interact with collagen (26,27) although this interaction is not well understood (28). The extracellular N-terminal regions of the  $\alpha$  and  $\beta$  integrin subunits are important in the bind-30 ing of ligands (29,30). The N-terminal region of the  $\alpha$ -subunits is composed of a seven-fold repeated sequence (12,31) containing FG and GAP consensus sequences. The repeats are predicted to fold into a β-propeller domain

15

25

30

(32) with the last three or four repeats containing putative divalent cation binding sites. The  $\alpha$ -integrin subunits  $\alpha$ 1,  $\alpha$ 2,  $\alpha$ D,  $\alpha$ E,  $\alpha$ L,  $\alpha$ M and  $\alpha$ X contain a ~200 amino acid inserted domain, the I-domain (A-domain), which shows similarity to sequences in von Willebrand factor, cartilage matrix protein and complement factors C2 and B (33,34). The I-domain is localized between the second and third FG-GAP repeats, it contains a metal ion-dependent adhesion site (MIDAS) and it is involved in binding of ligands (35-38).

Chondrocytes, the only type of cells in cartilage, express a number of different integrins including  $\alpha 1\beta 1$ ,  $\alpha 2\beta 1$ ,  $\alpha 3\beta 1$ ,  $\alpha 5\beta 1$ ,  $\alpha 6\beta 1$ ,  $\alpha v\beta 3$ , and  $\alpha v\beta 5$  (39-41). It has been shown that  $\alpha 1\beta 1$  and  $\alpha 2\beta 1$  mediate chondrocyte interactions with collagen type II (25) which is one of the major components in cartilage. It has also been shown that  $\alpha 2\beta 1$  is a receptor for the cartilage matrix protein chondroadherin (42).

# 20 SUMMARY OF THE INVENTION

The present invention relates to a novel collagen type II binding integrin, comprising a subunit  $\alpha 10$  in association with a subunit  $\beta$ , especially subunit  $\beta 1$ , but also other  $\beta$ -subunits may be contemplated. In preferred embodiments, this integrin has been isolated from human or bovine articular chondrocytes, and human chondrosarcoma cells.

The invention also encompasses integrin homologues of said integrin, isolated from other species, such as bovine integrin heterodimer comprising a subunit  $\alpha 10$  in association with a subunit  $\beta$ , preferably  $\beta 1$ , as well as homologues isolated from other types of human cells or from cells originating from other species.

The present invention relates in particular to a recombinant or isolated integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, and homologues and or fragments thereof having the

25

30

same biological activity.

The invention further relates to a process of producing a recombinant integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity, which process comprises the steps of

- a) isolating a polynucleotide comprising a nucleotide sequence coding for a integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity,
- b) constructing an expression vector comprising the isolated polynucleotide,
- c) transforming a host cell with said expression vector,
- d) culturing said transformed host cell in a culture medium under conditions suitable for expression of integrin subunit α10, or homologues or fragments thereof having similar biological activity, in said transformed host cell, and, optionally,
- e) isolating the integrin subunit  $\alpha 10$ , or homologues or fragments thereof having the same biological activity, from said transformed host cell or said culture medium.

The integrin subunit  $\alpha 10$ , or homologues or fragments thereof having the same biological activity, can also be provided by isolation from a cell in which they are naturally present.

The invention also relates to an isolated polynucleotide comprising a nucleotide coding for a integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, which polynucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or parts thereof.

The invention further relates to an isolated polynucleotide or oligonucleotide which hybridises to a DNA or RNA encoding an integrin subunit  $\alpha 10$ , having the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof, wherein said polyoligo-

15

20

25

30

nucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding the integrin subunit  $\alpha 1$ .

The invention relates in a further aspect to vectors comprising the above polynucleotides, and to cells containing said vectors and cells that have polynucleotides or oligonycleotides as shown in SEQ ID No. 1 or 2 integrated in their genome.

The invention also relates to binding entities having the capability of binding specifically to the integrin subunit  $\alpha 10$  or to homologues or fragments thereof, such as proteins, peptides, carbohydrates, lipids, natural ligands, polyclonal antibodies or monoclonal antibodies.

In a further aspect, the invention relates to a recombinant or isolated integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity.

In a preferred embodiment thereof, the subunit  $\beta$  is  $\beta 1.$ 

The invention also relates to a process of producing a recombinant integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, which process comprises the steps of

- a) isolating one polynucleotide comprising a nucleotide sequence coding for a subunit  $\alpha 10$  of an integrin heterodimer and, optionally, another polynucleotide comprising a nucleotide sequence coding for a subunit  $\beta$  of an integrin heterodimer, or for homologues or fragments thereof having similar biological activity,
- b) constructing an expression vector comprising said isolated polynucleotide coding for said subunit  $\alpha 10$  in combination with an expression vector comprising said isolated nucleotide coding for said subunit  $\beta$ ,

15

20

25

30

35

- c) transforming a host cell with said expression vectors,
- d) culturing said transformed host cell in a culture medium under conditions suitable for expression of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof similar biological activity, in said transformed host cell, and, optionally,
- e) isolating the integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof having the same biological activity, from said transformed host cell or said culture medium.

The integrin heterodimer, or homologues or fragments thereof having similar biological activity, can also be provided by isolation from a cell in which they are naturally present.

The invention further relates to a cell containing a first vector, said first vector comprising a polynucleotide coding for a subunit  $\alpha 10$  of an integrin heterodimer, or for homologues or parts thereof having similar biological activity, which polynucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof, and, optionally, a second vector, said second vector comprising a polynucleotide coding for a subunit  $\beta$  of an integrin heterodimer, or for homologues or fragments thereof.

In still another aspect, the invention relates to binding entities having the capability of binding specifically to the integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, preferably wherein the subunit  $\beta$  is  $\beta 1$ . Preferred binding entities are proteins, peptides, carbohydrates, lipids, natural ligands, polyclonal antibodies and monoclonal antibodies.

In a further aspect, the invention relates to a fragment of the integrin subunit  $\alpha 10$ , which fragment is a peptide chosen from the group comprising peptides of

10

15

20

25

30

35

the cytoplasmic domain, the I-domain and the spliced domain.

In one embodiment, said fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.

In another embodiment, said fragment comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEQ ID No. 1.

In a further embodiment, said fragment comprises the amino acid sequence from about amino acid No. 140 to about amino acid No. 337 in SEQ ID No. 1.

Another embodiment of the invention relates to a polynucleotide or oligonucleotide coding for a fragment of the human integrin subunit  $\alpha 10$ . In one embodiment this polynucleotide of oligonucleotide codes for a fragment which is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain. In further embodiments the polynucleotide or oligonucleotide codes for the fragments defined above.

The invention also relates to binding entities having the capability of binding specifically to a fragment of the integrin subunit  $\alpha 10$  as defined above.

The invention also relates to a process of using an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or a homologue or fragment of said integrin or subunit having similar biological activity, as a marker or target molecule of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.

In an embodiment of this process the fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.

In further embodiments of said process the fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or a fragment comprising the amino acid sequence from about amino acid No. 952 to

15

20

25

30

35

about amino acid No. 986 of SEQ ID No. 1, or a fragment comprising the amino acid sequence from about amino acid no. 140 to about amino acid no. 337 of SEQ ID no. 1.

The subunit  $\beta$  is preferably  $\beta 1$ . The cells are preferably chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

Said process may be used during pathological conditions involving said subunit  $\alpha 10$ , such as pathological conditions comprising damage of cartilage, or comprising trauma, rheumatoid arthritis and osteoarthritis.

Said process may be used for detecting the formation of cartilage during embryonal development, or for detecting physiological or therapeutic reparation of cartilage.

Said process may also be used for selection and analysis, or for sorting, isolating or purification of chondrocytes.

A further embodiment of said process is a process for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes.

A still further embodiment of said process is a process for in vitro studies of differentiation of chondrocytes.

The invention also comprises a process of using binding entities having the capability of binding specifically to an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, as markers or target molecules of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.

The fragment in said process may be a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain. In preferred embodiments said fragment is a peptide comprising the

15

20

amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or a fragment comprising the amino acid sequence from about amino acid No. 952 to about amino acid No. 986 of SEQ ID No. 1, or a fragment comprising the amino acid sequence from about amino acid No. 140 to about amino acid no. 337 of SEQ ID No. 1.

The process may also be used for detecting the presence of an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or of an integrin heterodimer comprising said subunit  $\alpha 10$ and a subunit  $\beta$ , or of homologues or fragments thereof having similar biological activity.

In a further embodiment said process is a process for determining the differentiation-state of cells during embryonic development, angiogenesis, or development of cancer.

In a still further embodiment this process is a process for detecting the presence of an integrin subunit  $\alpha 10$ , or of a homologue or fragment of said integrin subunit having similar biological activity, on cells, whereby a polynucleotide or oligonucleotide chosen from the group comprising a polynucleotide or oligonucleotide chosen from the nucelotide sequence shown in SEQ ID No. 1 is used as a marker under hybridisation conditions wherein said polynucleotide or oligonucleotide fails to 25 hybridise to a DNA or RNA encoding an integrin subunit lphal. Said cells may be chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts. Said integrin fragment may be a peptide chosen from the group comprising peptides 30 of the cytoplasmic domain, the I-domain and the spliced domain, such as a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or a fragment comprising the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEQ ID No. 1, or a fragment comprising the amino acid sequence from about amino acid No. 140 to about amino acid no. 337 of SEQ ID No. 1.

In a still further embodiment the process is a process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration or in therapeutic and physiological reparation of cartilage. The pathological conditions may be any pathological conditions involving the integrin subunit  $\alpha 10$ , such as rheumatoid arthritis, osteoarthrosis or cancer. The cells may be chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

The invention also relates to a process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration and in therapeutic and physiological reparation of cartilage, whereby a polynucleotide or oligonucleotide 15 chosen from the nucleotide sequence shown in SEQ ID No. 1 is used as a marker under hybridisation conditions wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit lphal. Embodiments of this aspect comprise a process, where-20 by said polynucleotide or oligonucleotide is a polynucleotide or oligonucleotide coding for a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain, such as a polynucleotide or oligonucleotide coding 25 for a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ, or comprising the amino acid sequence from about amino acid No. 952 to about amino acid no. 986 of SEQ ID No. 1, or the amino acid sequence from about amino acid No. 140 to about amino acid No. 337 30 of SEQ ID No. 1. Said pathological conditions may be any pathological conditions involving the integrin subunit  $\alpha$ 10, such as rheumatoid arthritis, osteoarthrosis or cancer, or atherosclerosis or inflammation. Said cells may be chosen from the group comprising chondrocytes, 35 smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

20

25

30

35

In a further aspect the invention relates to a pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$ and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule. An embodiment of said pharmaceutical composition is intended for use in stimulating, inhibiting or blocking the formation of cartilage, bone or blood vessels. A further embodiment comprises a pharmaceutical composition for use in preventing adhesion between tendon/ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue. 15

The invention is also related to a vaccine comprising as an active ingredient an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$ thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$ , or DNA or RNA coding for said integrin subunit alo.

A further aspect of the invention is related to the use of the integrin subunit  $\alpha 10$  as defined above as a marker or target in transplantation of cartilage or chondrocytes.

A still further aspect of the invention is related to a method of using binding entities having the capability of binding specifically to an integrin subunit  $\alpha 10$ comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, for promoting adhesion of chondrocytes and/or osteoblasts to surfaces of implants to stimulate osseointegration.

The invention is also related to the use of an integrin subunit  $\alpha 10$  or an integrin heterodimer comprising said subunit lpha 10 and a subunit  $oldsymbol{eta}$  as a target for anti-

15

20

30

adhesive drugs or molecules in tendon, ligament, skeletal muscle or other tissues where adhesion impairs the function of the tissue.

The invention also relates to a method of stimulating, inhibiting or blocking the formation of cartilage or bone, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit 10 α10 having similar biological activity, as a target molecule.

In another embodiment the invention is related to a method of preventing adhesion between tendon/ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using a integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.

The invention also relates to a method of stimulating extracellular matrix synthesis and repair by activation or blockage of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta \text{, or of the subunit }\alpha 10$ thereof, or of a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity.

In a further aspect the invention relates to a method of in vitro detecting the presence of integrin binding entities, comprising interaction of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, with a sample, thereby causing said integrin, subunit  $\alpha 10$ , or homologue or fragment thereof having similar biological activity, to modulate

10

15

the binding to its natural ligand or other integrin binding proteins present in said sample.

The invention also relates to a method of in vitro studying consequences of the interaction of a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, with an integrin binding entity and thereby initiate a cellular reaction. Said consequences may be measured as alterations in cellular functions.

A still further aspect of the inventions relates to a method of using DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof as a molecular target. In an embodiment of this aspect, a polynucleotide or oligonucleotide hybridises to the DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, whereby said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding en integrin subunit  $\alpha 1$ .

The invention also relates to a method of using a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, or a DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, as a marker or target molecule during angiogenesis.

BRIEF DESCRIPTION OF THE FIGURES

Fig.1 Affinity purification of the  $\alpha 10$  integrin subunit on collagen type II-Sepharose.

Fig. 2. Amino acid sequences of peptides from the bovine  $\alpha 10$  integrin subunit.

Fig. 3a. Affinitypurification and immunoprecipitation of the integrin subunit  $\alpha 10$  from bovine chondrocytes.

Fig. 3b. Affinitypurification and immunoprecipitation of the integrin subunit  $\alpha 10$  from human chondrocytes.

PCT/SE99/00544

25

35

- Fig. 3c. Affinitypurification and immunoprecipitation of the integrin subunit  $\alpha 10$  from human chondrosarcoma cells.
- Fig. 4. A 900 bp PCR-fragment corresponding to the bovine integrin subunit  $\alpha 10$ 
  - Fig. 5. Schematic map of the three overlapping  $\alpha 10$  clones.
  - Fig. 6. Nucleotide sequence and deduced amino acid sequence of the human  $\alpha 10$  integrin subunit.
- 10 Fig. 7. Northern blot of integrin α10 mRNA.
  - Fig. 8 Immunoprecipitation of the  $\alpha 10$  integrin subunit from human chondrocytes using antibodies against the cytoplasmic domain of  $\alpha 10$  (a). Western blot of the  $\alpha 10$  associated  $\beta$ -chain (b).
- 15 Fig. 9. Immunostaining of  $\alpha 10$  integrin in human articular cartilage.
  - Fig. 10 Immunostaining of  $\alpha 10$  integrin in 3 day mouse limb cartilage.
- Fig 11. Immunostaining of  $\alpha 10$  integrin in 13.5 day 20 mouse embryo.
  - Fig 12. Hybridisation of  $\alpha 10\ mRNA$  in various human tissues.
  - Fig. 13 Immunostaining of fascia around tendon (a), skeletal muscle (b) and heart valves (c) in 3 day mouse limb.
  - Fig. 14. PCR fragments corresponding to  $\alpha$ 10 integrin subunit from human chondrocytes, human endothelial cells, human fibroblasts and rat tendon.
- Fig 15. Partial genomic nucleotide sequence of the 30 human integrin subunit  $\alpha 10$ .
  - Fig 16. Upregulation of  $\alpha 10$  integrin subunit in chondrocytes cultured in alginate.
  - Fig 17. Immunoprecipitation of the  $\alpha 10$  integrin subunit from human smooth muscle cells

DETAILED DESCRIPTION OF THE INVENTION

The present invention demonstrate that human and

30

35

bovine chondrocytes express a novel, collagen type II-binding integrin in the  $\beta$ 1-family. An earlier study presented some evidence for that human chondrosarcoma cells also express this integrin (25). Immunoprecipitation experiments using antibodies against the integrin subunit  $\beta 1$  revealed that this novel  $\alpha$ -integrin subunit had an apparent molecular weight  $(M_{\rm r}$  ) of approximately 160 kDa under reducing conditions, and was slightly larger than the lpha2 integrin subunit. To isolate this  $\alpha$ -subunit collagen type II-binding proteins were affinity 10 purified from bovine chondrocytes. The chondrocyte lysate was first applied to a fibronectin-Sepharose precolumn and the flow through was then applied to a collagen type II-Sepharose column. A protein with  $M_{\rm r}$  of approximately 160 kD was specifically eluted with EDTA from the colla-15 gen column but not from the fibronectin column. The  $M_{\mathtt{r}}$  of this protein corresponded with the  $M_{\rm r}$  of the unidentified  $\beta$ 1-related integrin subunit. The 160 kD protein band was excised from the SDS-PAGE gel, digested with trypsin and the amino acid sequences of the isolated peptides were 20 analysed.

Primers corresponding to isolated peptides amplified a 900 bp PCR-fragment from bovine cDNA which was cloned, sequenced and used for screening of a human articular chondrocyte  $\lambda Z$ apII cDNA library to obtain the human integrin  $\alpha$ -subunit homologue. Two overlapping clones, hcl and hc2 were isolated, subcloned and sequenced. These clones contained 2/3 of the nucleotide sequence including the 3' end of the cDNA. A third clone which contained the 5'end of the  $\alpha$ 10 cDNA, was obtained using the RACE technique. Sequence analysis of the 160 kD protein sequence showed that it was a member of the integrin  $\alpha$ -subunit family and the protein was named  $\alpha$ 10.

The deduced amino acid sequence of  $\alpha 10$  was found to share the general structure of the integrin  $\alpha$ -subunits described in previously published reports (6-21). The large extracellular N-terminal part of  $\alpha 10$  contains a

30

35

seven-fold repeated sequence which was recently predicted to fold into a  $\beta$ -propeller domain (32). The integrin subunit  $\alpha 10$  contains three putative divalent cation-binding. sites (DxD/NxD/NxxxD) (53), a single spanning transmembrane domain and a short cytoplasmic domain. In contrast to most  $\alpha$ -integrin subunits the cytoplasmic domain of  $\alpha 10$ does not contain the conserved sequence KxGFF (R/K) R. The predicted amino acid sequence in lpha 10 is KLGFFAH. Several reports indicate that the integrin cytoplasmic domains are crucial in signal transduction (54) and that membrane-proximal regions of both  $\alpha-$  and  $\beta-$ integrin cytoplasmic domains are involved in modulating conformation and affinity state of integrins (55-57). It is suggested that the GFFKR motif in  $\alpha$ -chains are important for association of integrin subunits and for transport of the 15 integrin to the plasma membrane (58). The KxGFFKR domain has been shown to interact with the intracellular protein calreticulin (59) and interestingly, calreticulin-null embryonic stem cells are deficient in integrin-mediated cell adhesion (60). It is therefor possible that the 20 sequence KLGFFAH in alo have a key function in regulating the affinity between  $\alpha 10\beta 1$  and matrix proteins.

Integrin  $\alpha$  subunits are known to share an overall identity of 20-40% (61). Sequence analysis showed that the  $\alpha$ 10 subunit is most closely related to the I-domain containing  $\alpha$ -subunits with the highest identity to  $\alpha$ 1 (37%) and  $\alpha$ 2 (35%). The integrins  $\alpha$ 1 $\beta$ 1 and  $\alpha$ 2 $\beta$ 1 are known receptors for both collagens and laminins (24;62;63) and we have also recently demonstrated that  $\alpha$ 2 $\beta$ 1 interacts with the cartilage matrix protein chondroadherin (42). Since  $\alpha$ 10 $\beta$ 1 was isolated on a collagen type II-Sepharose, we know that collagen type II is a ligand for  $\alpha$ 10 $\beta$ 1. We have also shown by affinity purification experiments that  $\alpha$ 10 $\beta$ 1 interacts with collagen type I but it remains to be seen whether laminin or chondroadherin are also ligands for this integrin.

15

20

25

30

35

The  $\alpha 10$  associated  $\beta$ -chain migrated as the  $\beta 1$  integrin subunit both under reducing and non-reducing conditions. To verify that the  $\alpha 10$  associated  $\beta$ -chain indeed is  $\beta 1$ , chondrocyte lysates were immunoprecipitated with antibodies against  $\alpha 10$  or  $\beta 1$  followed by Western blot using antibodies against the  $\beta 1$ -subunit. These results clearly demonstrated that  $\alpha 10$  is a member of the  $\beta 1$ -integrin family. However, the possibility that  $\alpha 10$  combine also with other  $\beta$ -chains can not be excluded..

A polyclonal peptide antibody raised against the cytoplasmic domain of  $\alpha 10$  precipitated two protein bands with  $\text{M}_{\text{r}}$  of approximately 160 kD ( $\alpha 10)$  and 125 kD ( $\beta 1)$ under reducing conditions. Immunohistochemistry using the  $\alpha$ 10-antibody showed staining of the chondrocytes in tissue sections of human articular cartilage. The antibody staining was clearly specific since preincubation of the antibody with the  $\alpha 10$ -peptide completely abolished the staining. Immunohistochemical staining of mouse limb sections from embryonic tissue demonstrated that  $\alpha 10$  is upregulated during condensation of the mesenchyme. This indicate that the integrin subunit  $\alpha 10$  is important during the formation of cartilage. In 3 day old mice  $\alpha 10$ was found to be the dominating collagen binding integrin subunit which point to that  $\alpha 10$  has a key function in maintaining normal cartilage functions.

Expression studies on the protein and mRNA level show that the distribution of  $\alpha 10$  is rather restrictive. Immunohistochemistry analyses have shown that  $\alpha 10$  integrin subunit is mainly expressed in cartilage but it is also found in perichondrium, periosteum, ossification groove of Ranvier, in fascia surrounding tendon and skeletal muscle and in the tendon-like structures in the heart valves. This distribution point to that  $\alpha 10$  integrin subunit is present also on fibroblasts and osteoblasts. PCR amplification of cDNA from different cell types revealed the presence of an alternatively spliced  $\alpha 10$  integrin subunit. This spliced  $\alpha 10$  was domi-

nating in fibroblasts which suggests that  $\alpha 10$  in fibroblasts may have a different function compared to  $\alpha 10$  present on chondrocytes.

Expression of the integrin subunit  $\alpha 10$  was found to decrease when chondrocytes were cultured in monolayer. In contrast, the expression of  $\alpha 10$  was found to increase when the cells were cultured in alginate beads. Since the latter culturing model is known to preserve the phenotype of chondrocytes the results suggest that  $\alpha 10$  can function as marker for a differentiated chondrocyte.

Adhesion between tendon/ligaments and the surrounding tissue is a well-known problem after infection, injury and after surgical intervention. Adhesion between tendon and tendon sheets impairs the gliding function and cause considerable problems especially during healing of tendons in e.g. the hand and fingers leading to functional incapacity. The localisation of the  $\alpha 10$  integrin subunit in the fascia of tendon and skeletal muscle makes  $\alpha 10$  a possible target for drugs and molecules with antiadhesive properties that could prevent impairment of the function of tendon/ligament. The integrin subunit  $\alpha 10$  can also be a target for drugs or molecules with antiadhesive properties in other tissues where adhesion is a problem.

25

35

20

10

15

#### EXAMPLES

#### Example 1

Affinity purification of the  $\alpha 10$  integrin subunit on collagen type II-Sepharose.

# Materials and Methods

Bovine chondrocytes, human chondrocytes or human chondrosarcoma cells were isolated as described earlier [Holmvall et al, Exp Cell Res, 221, 496-503 (1995), Camper et al, JBC, 273, 20383-20389 (1998)]. A Triton X-100 lysate of bovine chondrocytes was applied to a fibronectin-Sepharose precolumn followed by a collagen

type II-Sepharose column and the integrin subunit α10 was eluted from the collagen type II-column by EDTA (Camper et al, JBC, 273, 20383-20389 (1998). The eluted proteins were precipitated by methanol/chloroform, separated by SDS-PAGE under reducing conditions and stained with Coomassie blue. (Camper et al, JBC, 273, 20383-20389 (1998). Peptides from the α10 protein band were isolated by in-gel digestion with a trypsin and phase liquid chromatography and sequenced by Edman degradation (Camper et al, JBC, 273, 20383-20389 (1998). Results

Fig 1 shows EDTA-eluted proteins from the fibronectin-Sepharose (A), flow-through from the collagen type II-Sepharose column (B) and EDTA-eluted proteins from the collagen type II-Sepharose (C). The  $\alpha 10$  integrin subunit 15 (160 kDa) which was specifically eluted from the collagen type II column is indicated with an arrow. Figure 2 shows the amino acid sequences of six peptides that were isolated from the bovine integrin subunit lpha 10. Figures 3 a, b, and c show that the  $\alpha 10$  integrin subunit is present on bovine chondrocytes (3a), human chondrocytes (3b) and human chondrosarcoma cells (3c). The affinity for collagen type II, the coprecipitation with  $\beta 1$ -integrin subunit and the molecular weight of 160 kDa under reducing conditions identify the  $\alpha 10$  integrin subunit on the different cells. These results show that  $\alpha 10$  can be isolated from chondrocytes and from chondrosarcoma cells.

Example 2

30

35

Amplification of PCR fragment corresponding to bovine  $\alpha 10$  integrin subunit. Materials and methods

The degenerate primers GAY AAY ACI GCI CAR AC (DNTAQT, forward) and TIA TIS WRT GRT GIG GYT (EPHHSI, reverse) were used in PCR (Camper et al, JBC, 273, 20383-20389 (1998) to amplify the nucleotide sequence corresponding to the bovine peptide 1 (Figure 2). A 900 bp

PCR-fragment was then amplified from bovine cDNA using an internal specific primer TCA GCC TAC ATT CAG TAT (SAYIQY, forward) corresponding to the cloned nucleotide sequence of peptide 1 together with the degenerate primer ICK RTC CCA RTG ICC IGG (PGHWDR, reverse) corresponding to the bovine peptide 2 (Figure2). Mixed bases were used in positions that were twofold degenerate and inosines were used in positions that are three- or fourfold degenerate. mRNA isolation and cDNA synthesis was done as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)). The purified fragment was cloned, purified and sequenced as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)).

Results

The nucleotide sequence of peptide 1 (Figure 2)
was obtained by PCR-amplification, cloning and sequencing of bovine cDNA. From this nucleotide sequence an
exact primer was designed and applied in PCR-amplification with degenerate primers corresponding to peptides
20 2-6 (Figure 2). Primers corresponding to peptides 1
and 2 amplified a 900 bp PCR-fragment from bovine cDNA
(Figure 4).

#### Example 3

25

30

35

Cloning and sequence analysis of the human  $\alpha 10$  integrin subunit

Material and methods

The cloned 900bp PCR-fragment, corresponding to bovine alo-integrin, was digoxigenin-labelled according to the DIG DNA labelling kit (Boehringer Mannheim) and used as a probe for screening of a human articular chondrocyte \(\lambda\text{ZapII}\) cDNA library (provided by Michael Bayliss, The Royal Veterinary Basic Sciences, London, UK) (52). Positive clones containing the pBluescript SK+ plasmid with the cDNA insert were rescued from the ZAP vector by in vivo excision as described in the ZAP-cDNA® synthesis kit (Stratagene). Selected plasmids were purified and

PCT/SE99/00544

sequenced as described earlier (Camper et al, JBC, 273, 20383-20389 (1998)) using T3, T7 and internal specific primers. To obtain cDNA that encoded the 5' end of  $\alpha$ 10 we designed the primer AAC TCG TCT TCC AGT GCC ATT CGT GGG (reverse; residue 1254-1280 in  $\alpha$ 10 cDNA) and used it for rapid amplification of the cDNA 5' end (RACE) as described in the Marathon CDNA Amplification kit (Clontech INC., Palo Alto, CA).

Results

Two overlapping clones, hcl and hc2 (Figure 5), were 10 isolated, subcloned and sequenced. These clones contained 2/3 of the nucleotide sequence including the 3' end of the cDNA. A third clone (racel; Figure 5), which contained the 5'end of the  $\alpha 10$  cDNA, was obtained using the RACE technique. From these three overlapping clones of 15  $\alpha$ 10 cDNA, 3884 nucleotides were sequenced The nucleotide sequence and deduced amino acid sequence is shown in Figure 6. The sequence contains a 3504-nucleotide open reading frame that is predicted to encode a 1167 amino acid mature protein. The signal peptide cleavage site is 20 marked with an arrow, human homologues to bovine peptide sequences are underlined and the I-domain is boxed. Metal ion binding sites are indicated with a broken underline, potential N-glycosylation sites are indicated by an asterisk and the putative transmembrane domain is double 25 underlined. The normally conserved cytoplasmic sequence is indicated by a dot and dashed broken underline.

Sequence analysis demonstrate that  $\alpha 10$  is a member of the integrin  $\alpha\text{-subunit}$  family.

30

35

Example 4

Identification of a clone containing a splice variant of  $\alpha 10$ 

One clone which was isolated from the human chondrocyte library (see Example 3) contained a sequence that was identical to the sequence of  $\alpha 10$  integrin subunit except that the nucleotides between nt positions

2942 and 3055 were deleted. The splice variant of  $\alpha 10$  was verified in PCR experiment using primers flanking the splice region (see figure 14).

#### 5 Example 5

30

35

Identification of  $\alpha 10$  integrin subunit by Northern blot

Material and methods

Bovine chondrocyte mRNA was purified using a QuickPrep®Micro mRNA Purification Kit (Pharmacia Biotech, 10 Uppsala, Sweden), separated on a 1% agarose-formaldehyde gel, transferred to nylon membranes and immobilised by UV crosslinking. cDNA-probes were 32P-labelled with Random Primed DNA Labeling Kit (Boehringer Mannheim). Filters were prehybridised for 2-4 hours at 42°C in 5x SSE, 15 5x Denharts solution, 0.1 % SDS, 50 μg/ml salmon sperm DNA and 50% formamide and then hybridised over night at 42 °C with the same solution containing the specific probe (0.5-1 x 106 cpm/ml). Specifically bound cDNAprobes were analysed using the phosphoimager system (Fuji). Filters were stripped by washing in 0.1% SDS, for 1 hour at 80°C prior to re-probing. The α10-integrin cDNA-probe was isolated from the racel-containing plasmid using the restriction enzymes BamHI (GIBCO BRL) and NcoI (Boehringer Mannheim). The rat  $\beta 1$ -integrin cDNA probe was 25 a kind gift from Staffan Johansson, Uppsala, Sweden. Results

Northern blot analysis of mRNA from bovine chondrocytes showed that a human  $\alpha 10$  cDNA-probe hybridised with a single mRNA of approximately 5.4 kb (Figure 7). As a comparison, a cDNA-probe corresponding to the integrin subunit  $\alpha 1$  was used. This cDNA-probe hybridised a mRNA-band of approximately 3.5 kb on the same filter. These results show that a cDNA-probe against  $\alpha 10$  can be used to identify the  $\alpha 10$  integrin subunit on the mRNA level.

Example 6

Preparation of antibodies against the integrin subunit  $\alpha 10$ 

A peptide corresponding to part of the  $\alpha 10$  cytoplasmic domain, Ckkipeeekreekle (see figure 6) was synthesised and conjugated to keyhole limpet hemocyanin (KLH). Rabbits were immunised with the peptide-KLH conjugate to generate antiserum against the integrin subunit  $\alpha 10$ . Antibodies recognising  $\alpha 10$  were affinity purified on an peptide-coupled column (Innovagen AB).

#### Example 7

10

Immunoprecipitation of the integrin subunit  $\alpha 10$  from chondrocytes

15 Material and methods

Human chondrocytes were 125I-labelled, lyzed with Triton X-100 and immunoprecipitated as earlier described (Holmvall et al, Exp Cell Res, 221, 496-503 (1995), Camper et al, JBC, 273, 20383-20389 (1998)). Triton X-100 lysates of 125I-labeled human chondrocytes were immuno-20 precipitated with polyclonal antibodies against the integrin subunits  $\beta$ 1,  $\alpha$ 1,  $\alpha$ 2,  $\alpha$ 3 or  $\alpha$ 10. The immunoprecipitated proteins were separated by SDS-PAGE (4-12%) under non-reducing conditions and visualised using a phosphoimager. Triton X-100 lysates of human chondrocytes immu-25 noprecipitated with  $\alpha 10$  or  $\beta 1$  were separated by SDS-PAGE (8%) under non-reducing conditions and analysed by Western blot using the polyclonal  $\beta1$  antibody and chemiluminescent detection as described in Camper et al, JBC, 273, 20383-20389 (1998). 30

#### Results

The polyclonal peptide antibody, raised against the cytoplasmic domain of  $\alpha 10$ , precipitated two protein bands with Mr of approximately 160 kD ( $\alpha 10$ ) and 125 kD ( $\beta 1$ ) under reducing conditions. The  $\alpha 10$  associated  $\beta$ -chain migrated as the  $\beta 1$  integrin subunit (Figure 8a). To verify that the  $\alpha 10$  associated  $\beta$ -chain in chondrocytes

indeed is  $\beta 1$ , chondrocyte lysates were immunoprecipitated with antibodies against  $\alpha 10$  orb  $\beta 1$  followed by Western blot using antibodies against the  $\beta 1$ -subunit (Figure 8b). These results clearly demonstrated that  $\alpha 10$  is a member of the  $\beta 1$ -integrin family. However, the results do not exclude the possibility that  $\alpha 10$  can associate with other  $\beta$ -chains in other situations.

#### Example 8

10

25

30

35

Immunohistochemical staining of the integrin subunit  $\alpha 10$  in human and mouse cartilage Material and methods

Frozen sections of adult cartilage (trochlear groove) obtained during surgery (provided by Anders

Lindahl, Salgrenska Hospital, Gothenburg, Sweden and frozen sections from of 3 day old mouse limb were fixed and prepared for immunohistochemistry as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)). Expression of α10 integrin subunit was analysed using the polyclonal antibody against the cytoplasmic domain as a primary antibody (see Example 6) and a secondary antibody conjugated to peroxidase.

Results

Figures 9 show immunostaining of human adult articular cartilage.

The  $\alpha 10$ -antibody recognising the cytoplasmic domain of  $\alpha 10$  stained the chondrocytes in tissue sections of human articular cartilage (A). The staining was depleted when the antibody was preincubated with the  $\alpha 10$ - peptide (B). A control antibody recognising the  $\alpha 9$  integrin subunit did not bind to the chondrocyte (C).

Figures 10 shows that the  $\alpha$ 10 antibody stain the majority of chondrocytes in the growing bone anlage (a and b). The  $\alpha$ 10 antibody also recognised cells in the ossification groove of Ranvier (b), especially the osteoblast in the bone bark which are lining the cartilage in the metaphys are highly positive for  $\alpha$ 10. The

cells in the ossification groove of Ranvier are believed to be important for the growth in diameter of the bone. The integrin subunit  $\alpha 10$  is also highly expressed in perichondrium and periosteum. Cell in these tissues are likely important in the repair of the cartilage tissue. The described localisation of the integrin subunit  $\alpha 10$  suggest that this integrin is important for the function of the cartilage tissue.

# 10 Example 9

Immunohistochemical staining of the integrin subunit  $\alpha 10\ during$  mouse development Material and methods

Frozen sections from mouse embryos (13.5 days) were investigated for expression of α10 by immunhistochemistry as described in Camper et al, JBC, 273, 20383-20389 (1998). Expression of α10 integrin subunit was analysed using the polyclonal antibody against the cytoplasmic domain as a primary antibody (see Example 6) and a secondary antibody conjugated to peroxidase. The embryo sections were also investigated for expression of integrin subunit α1 (monoclonal antibody from Pharmingen) and collagen type II (monoclonal antibody, kind gift from Dr John Mo, Lund University, Sweden).

# 25 Results

30

35

Figure 11 show that  $\alpha 10$  integrin subunit is unregulated in the limb when the mesenchymal cells undergo condensation to form cartilage (a). Especially the edge of the newly formed cartilage has high expression of  $\alpha 10$ . The formation of cartilage is verified by the high expression of the cartilage specific collage type II (b). The control antibody against  $\alpha 1$  integrin subunit showed only weak expression on the cartilage (c). In other experiments expression of  $\alpha 10$  was found in all cartilage containing tissues in the 3 day old mouse including limbs, ribs and vertebrae. The upregulation of  $\alpha 10$  during formation of cartilage suggest that this integrin subunit is

important both in the development of cartilage and bone and in the repair of damaged cartilage tissue.

# Example 10

5

20

mRNA expression of  $\alpha 10$  in tissues other than articular cartilage

Material and methods

Expression of α10 integrin subunit was examined on the mRNA level in different human tissues. A Northern dot blot with immobilised mRNA from the listed tissues in Figure 12 was hybridised with an α10 integrin cDNA probe isolated from the race 1-containing plasmid using the restriction enzymes BamH1 and Ncol. The degree of hybridisation was analysed using a phospho imager. The following symbols denote mRNA level in increasing order: -, +, +++, ++++.

#### Results

Analysis of the hybridised mRNA showed that  $\alpha 10$  was expressed in aorta, trachea, spinal cord, heart, lung, and kidney (Figure 12). All other tissues appeared negative for  $\alpha 10$  expression. These results point to a restricted distribution of the  $\alpha 10$  integrin subunit.

# Example 11

Immunohistochemical staining of α10 in fascia around tendon and skeletal muscle and in tendon structures in heart valves.

# Materials and methods

Frozen sections of adult cartilage (trochlear groove) obtained during surgery (provided by Anders Lindahl, Salgrenska Hospital, Gothenburg, Sweden and frozen sections from of 3 day old mouse limb were fixed and prepared for immunohistochemistry as earlier described (Camper et al, JBC, 273, 20383-20389 (1998)). Expression of α10 integrin subunit was analysed using the polyclonal antibody against the cytoplasmic domain as a pri-

mary antibody (see Example 6) and a secondary antibody conjugated to peroxidase. Results

As shown in figures 13 expression of  $\alpha 10$  was found in the fascia surrounding tendon (a) and skeletal muscle 5 (b) and in the tendon structures in the heart valves (c). This localisation suggest that  $\alpha 10$  can bind to other matrix molecules in addition to the cartilage specific collagen type II. The localisation of the integrin  $\alpha 10$  on the surface of tendons indicate that  $\alpha 10$  can be involved 10 in unwanted adhesion that often occurs between tendon/ ligaments and the surrounding tissue after infection, injury or after surgery.

#### Example 12 15

25

mRNA expression of all integrin subuhit in chondrocytes, endothelial cells and fibroblasts. Material and methods

Isolation of mRNA, synthesis of cDNA and PCR amplification was done as earlier described (Camper et al, 20 JBC, 273, 20383-20389 (1998)). Results

Figure 14 shows PCR amplification of α10 cDNA from human articular chondrocytes (lanes A6 and B1), human umbilical vein endothelial cells (lane A2), human fibroblasts (lane A4) and rat tendon (Fig 14b, lane B2). Lanes 1, 3, and 5 in figure: 14 A show amplified fragments corresponding to the integrin subunit  $\alpha 2$  in endothelial cells, fibroblasts and chondrocytes, respectively. cDNAprimers corresponding to the  $\alpha 10$  sequence positions nt 2919-2943 (forward) and nt 3554-3578 (reverse) (see Figure 6) were used to amplify  $\alpha 10$  cDNA from the different cells. The figure shows that  $\alpha 10$  was amplified in all three cell types. Two fragments of  $\alpha 10$  was amplified which represent the intact form of  $\alpha 10$  (larger fragment) 35 and a splice variant (smaller fragment). The larger fragment was dominating in chondrocytes while the smaller fragment was more pronounced in tendon (B2).

# Example 13

Construction of  $\alpha 10$  mammalian expression vector. 5 The full length protein coding sequence of lpha 10 (combined from 3 clones, see figure 6) was inserted into the mammalian expression vector, pcDNA3.1/Zeo (Invitrogen). The vector contains SV40 promoter and Zeosin selection sequence. The  $\alpha 10$  containing expression vector was trans-10 fected into cells that express the eta 1-integrin subunit but lack expression of the  $\alpha 10$  subunit. Expression of the  $\alpha 10$  integrin subunit on the cell surface can be analysed by immunoprecipitation and/or flow cytometry using antibodies specific for  $\alpha 10$ . The ligand binding capacity and 15 the function of the inserted  $\alpha 10$  integrin' subunit can be demonstrated in cell adhesion experiment and in signalling experiments.

### 20 Example 14

25

Construction of mammalian expression vector containing a splice variant of  $\alpha 10\,.$ 

The full length protein coding sequence of the splice variant of  $\alpha 10$  (nt 2942-nt3055 deleted) was inserted into the mammalian expression vector pcDNA3 (see Example 13). Expression and function of the splice variant can be analysed as described in example 13 and compared with the intact  $\alpha 10$  integrin subunit.

## 30 Example 15

Partial isolation and characterisation of the  $\alpha 10\,$  integrin genomic DNA Material and methods

Human α10 cDNA, isolated from the racel-containing
35 plasmid using the restriction enzymes BamHI (GIBCO BRL)
and NcoI (Boehringer Mannheim), was <sup>32</sup>P-labelled and used
as a probe for screening of a mouse 129 cosmid library

(provided by Reinhard Fässler, Lund University). Positive clones were isolated and subcloned. Selected plasmids were purified and sequenced as described earlier (Camper et al, JBC, 273, 20383-20389 (1998)) using T3, T7 and internal specific primers. Primers corresponding to mouse genomic DNA were then constructed and used in PCR to amplify and identify the genomic sequence of  $\alpha$ 10 from the cosmid clones.

# Results

Figure 15 shows 7958 nt of the  $\alpha$ 10 gene. This partial genomic DNA sequence of  $\alpha$ 10 integrin contains 8 exons, and a Kozak sequence. The mouse genomic  $\alpha$ 10 sequence was used to generate a targeting vector for knockout experiments.

15

10

# Example 16

Upregulation of  $\alpha 10$  integrin subunit in chondrocytes cultured in alginate beads

# Material and methods:

Human chondrocytes cultured in monolayer for 2 weeks were detached with trypsin-EDTA and introduced into alginate beads. Chondrocytes cultured in alginate are known to preserve their phenotype while chondrocytes cultured in monolayer are dedifferentiated. After 11 days chondrocytes cultured either in alginate or on monolayer were isolated and surface labelled with <sup>125</sup>I. The α10 integrin subunit was then immunoprecipitated with polyclonal antibodies recognising the cytoplasmic domain of α10 (see Example 6 and Camper et al, JBC, 273, 20383-20389 (1998)).

#### Results

35

As shown in figure 16 chondrocytes cultured in alginate beads (lanes 3 and 4) upregulated their protein expression of  $\alpha 10\beta 1$ . This was in contrast to chondrocytes cultured in monolayer (lanes 1 and 2) which had a very low expression of  $\alpha 10\beta 1$ . Immunoprecipitation with ab control antibody is shown in lanes 1 and 3.It is known that

15

30

35

chondrocytes preserve their cartilage specific matrixproduction in alginate cultures but not in monolayer culture which point to that alginate preserve the phenotype of chondrocytes. These results support that  $\alpha 10$  integrin subunit can be used as a marker for differentiated chondrocytes.

#### Example 17

Immunoprecipitation of the  $\alpha 10$  integrin subunit from 10 human smooth muscle cells.

# Material and methods

Human smooth muscle cells were isolated from human aorta. After one week in culture the cells were  $^{125}I-$  labelled, lysed and immunoprecipitated with antibodies against the integrin subunit  $\beta1$  (lane 1),  $\alpha1$  (lane 2),  $\alpha2$  (lane 3),  $\alpha10$  (lane 4),  $\alpha3$  (lane 5), control (lane 6) (Figure 17). The experiment was done as described in Example 7.

#### Results

The  $\alpha 10$  antibody precipitated two bands from the smooth muscle cells corresponding to the  $\alpha 10$  and the  $\beta 1$  integrin subunit (Fig. 17).

#### Example 18

Construction of bacterial expression vector containing sequence for  $\alpha 10$  splice region.

A plasmid for intracellular expression in E. coli of the alternatively spliced region (amino acid pos. 952-986, SEQ. ID 1) was constructed as described. The alternatively spliced region were back-translated using the E. coli high frequency codon table, creating a cDNA sequence of 96% identity with the original sequence (SEQ. ID 1 nucleotide pos 2940-3044). Using sequence overlap extension (Horton et al., Biotechniques 8:528, 1990) primer  $\alpha$ 10pfor (tab. I) and  $\alpha$ 10prev (tab. I) was used to generate a double stranded fragment encoding the  $\alpha$ 10 amino acid sequence. This fragment was used as a PCR

template with primers  $\alpha 10 pfor2$  (tab. I) and  $\alpha 10 prev2$  (tab. I) in order to generate restriction enzyme site for sub-cloning in a pET vector containing the Z-domain of staphylococcal protein A, creating a fusion of the  $\alpha 10$  spliced region with the amino terminal of the Z-domain with trombin cleavage site residing in-between. The fragment generated in the second PCR reaction is shown (SEQ ID No. 3) also indicating the unique restriction enzymes used for sub-cloning in the expression vector.

Table I

a10pfor	5'- GTTCAGAACCTGGGTTGCTACGTTGTTTCCGGTCTGATCATCTCCGC TCTGCTGCCGGCTGT-3'				
α10pfor2	5'-GGGGCATATGGTTCAGAACCTGGGTTGCTACGTTG-3'				
α10prev	5'- GATAACCTGGGACAAGCTTAGGAAGTAGTTACCACCGTGAGCAACAG CCGGCAGCAGAGCGGA-3'				
α10prev2	5'- GGGGGGATCCGCGCGCACCAGGCCGCTGATAACCTGGGACAAGCTT AGGAAGT-3'				

15

20

#### REFERENCES

- 1. Springer, T.A. (1990) Nature 346, 425-434
- 2. Ruoslahti, E. (1991) J.Clin.Invest. 87, 1-5
- 3. Hynes, R.O. (1992) Cell 69, 11-25
- 5 4. Hemler, M.E. (1988) Immunol. Today 9, 109-113
  - 5. Yamada, K.M. (1991) J.Biol.Chem. 266, 12809-12812
  - Palmer, E.L., Ruegg, C., Ferrando, R., Pytela, R., and Sheppard, D. (1993) J. Cell Biol. 123, 1289-1297
  - 7. Takada, Y., Elices, M.J., Crouse, C., and Hemler, M.E. (1989) EMBO J. 8, 1361-1368
  - Poncz, M., Eisman, R., Heidenreich, R., Silver, S.M., Vilaire, G., Surrey, S., Schwartz, E., and Bennett, J.S. (1987) J.Biol.Chem. 262, 8476-8482
  - 9. Larson, R.S., Corbi, A.L., Berman, L., and Springer, T. (1989) J.Cell Biol. 108, 703-712
  - 10. Corbi, A.L., Kishimoto, T.K., Miller, L.J., and Springer, T.A. (1988) J.Biol.Chem. 263, 12403-12411
  - 11. Argraves, W.S., Suzuki, S., Arai, H., Thompson, K., Pierschbacher, M.D., and Ruoslahti, E. (1987) J.Cell Biol. 105, 1183-1190
  - 12. Corbi, A.L., Miller, L.J., O'Connor, K., Larson, R.S., and Springer, T.A. (1987) EMBO J. 6, 4023-4028
  - 13. Briesewitz, R., Epstein, M.R., and Marcantonio, E.E. (1993) J.Biol.Chem. 268, 2989-2996
- 25 14. Ziober, B.L., Vu, M.P., Waleh, N., Crawford, J., Lin, C.S., and Kramer, R.H. (1993) J.Biol.Chem. 268, 26773-26783
  - 15. Hogervorst, F., Kuikman, I., van Kessel, A.G., and Sonnenberg, A. (1991) Eur.J.Biochem. 199, 425-433
- 30 16. Takada, Y. and Hemler, M.E. (1989) J.Cell Biol. 109, 397-407
  - 17. Takada, Y., Murphy, E., Pil, P., Chen, C., Ginsberg, M.H., and Hemler, M.E. (1991) J.Cell Biol. 115, 257-266
- 35 18. Van der Vieren, M., Le Trong, H., Wood, C.L., Moore, P.F., St.John, T., Staunton, D.E., and Gallatin, W.M. (1995) Immunity. 3, 683-690

- Schnapp, L.M., Breuss, J.M., Ramos, D.M., Sheppard,
   D., and Pytela, R. (1995) J.Cell Sci. 108, 537-544
- 20. Shaw, S.K., Cepek, K.L., Murphy, E.A., Russell, G.J., Brenner, M.B., and Parker, C.M. (1994) J.Biol.Chem. 269, 6016-6025
- 21. Suzuki, S., Argraves, W.S., Arai, H., Languino, L.R., Pierschbacher, M.D., and Ruoslahti, E. (1987) J.Biol.Chem. 262, 14080-14085
- Ignatius, M.J., Large, T.H., Houde, M., Tawil, J.W.,
  Barton, A., Esch, F., Carbonetto, S., and Reichardt,
  L.F. (1990) J.Cell Biol. 111, 709-720
  - 23. Gullberg, D., Gehlsen, K.R., Turner, D.C., Åhlén, K., Zijenah, L.S., Barnes, M.J., and Rubin, K. (1992) EMBO J. 11, 3865-3873
- 15 24. Staaz, W.D., Rajpara, S.M., Wayner, E.A., Carter,
  W.G., and Santoro, S.A. (1989) J.Cell Biol. 108,
  1917-1924
  - 25. Holmvall, K., Camper, L., Johansson, S., Rubin, K., Kimura, J.H., and Lundgren-Åkerlund, E. (1995)
- 20 Exp.Cell Res. 221, 496-503
  - 26. Forsberg, E., Ek, B., Engström, Å., and Johansson, S. (1994) Exp.Cell Res. 213, 183-190
  - 27. Wayner, E.A. and Carter, W.G. (1987) J.Cell Biol. 105, 1873-1884
- 25 28. Weitzman, J.B., Pasqualini, R., Takada, Y., and Hemler, M.E. (1993) J.Biol.Chem. 268, 8651-8657
  - 29. Elices, M.J. and Hemler, M.E. (1989) Proc.Natl.Acad.Sci.U.S.A. 86, 9906-9910
- 30. Languino, L.R., Colella, S., Zanetti, A., Andrieux,
  30 A., Ryckewaert, J.J., Charon, M.H., Marchisio, P.C.,
  Plow, E.F., Ginsberg, M.H., Marguerie, G., and et al
  (1989) Blood 73, 734-742
  - 31. Tuckwell, D.S., Humphries, M.J., and Brass, A. (1994) Cell Adhes.Commun. 2, 385-402
- 35 32. Springer, T.A. (1997) Proc.Natl.Acad.Sci.U.S.A. 94, 65-72

- 33. Colombatti, A., Bonaldo, P., and Doliana, R. (1993)
  Matrix 13, 297-306
- 34. Lee, C.H., Bradley, G., and Ling, V. (1995) Cell Growth Differ. 6, 347-354
- 5 35. Calderwood, D.A., Tuckwell, D.S., and Humphries, M.J. (1995) Biochem.Soc.Trans. 23, 504S
  - 36. Kern, A., Eble, J., Golbik, R., and Kuhn, K. (1993) Eur.J.Biochem. 215, 151-159
- 37. Tuckwell, D.S., Reid, K.B., Barnes, M.J., and Humphries, M.J. (1996) Eur.J.Biochem. 241, 732-739
  - 38. Kamata, T. and Takada, Y. (1994) J.Biol.Chem. 269, 26006-26010
  - 39. Dürr, J., Goodman, S., Potocnik, A., von der Mark, H., and von der Mark, K. (1993) Exp.Cell Res. 207, 235-244
  - 40. Salter, D.M., Hughes, D.E., Simpson, R., and Gardner, D.L. (1992) Br.J.Rheumatol. 31, 231-234
  - 41. Woods, V.L.J., Schreck, P.J., Gesink, D.S., Pacheco, H.O., Amiel, D., Akeson, W.H., and Lotz, M. (1994)
    Arthritis Rheum. 37, 537-544
  - 42. Camper, L., Heinegård, D., and Lundgren-Åkerlund, E. (1997) J.Cell Biol 138, 1159-1167
  - 43. Hemler, M.E., Sanchez Madrid, F., Flotte, T.J., Krensky, A.M., Burakoff, S.J., Bhan, A.K., Springer,
- T.A., and Strominger, J.L. (1984) J.Immunol. 132, 3011-3018
  - 44. Bottger, B.A., Hedin, U., Johansson, S., and Thyberg, J. (1989) Differentiation. 41, 158-167
- 45. Sommarin, Y. and Heinegård, D. (1983) Biochem.J. 214, 777-784
  - 46. Häuselmann, H.J., Aydelotte, M.B., Schumacher, B.L., Kuettner, K.E., Gitelis, S.H., and Thonar, E.J.M.A. (1992) Matrix 12, 116-129
  - 47. Miller, E.J. (1972) Biochemistry 11, 4903-4909
- 35 48. Wessel, D. and Flugge, U.I. (1984) Anal.Biochem. 138, 141-143

- 49. Blobel, G. and Dobberstein, B. (1975) J.Cell Biol. 67, 835-851
- 50. Hellman, U. (1997) in Protein structure analysis.

  Preparation, characterization, and microsequencing
  (Kamp, R.M., Choli-Papadopoulou, T., and WittmannLiebold, B., eds) pp. 97-104, Spriner-Verlag,
  Heidelberg
  - 51. Charles, I.G., Palmer, R.M., Hickery, M.S., Bayliss, M.T., Chubb, A.P., Hall, V.S., Moss, D.W., and Moncada, S. (1993) Proc.Natl.Acad.Sci.U.S.A. 90, 11419-11423
  - 52. Tuckwell, D.S., Brass, A., and Humphries, M.J. (1992) Biochem.J. 285, 325-331
- 53. Dedhar, S. and Hannigan, G.E. (1996) Curr.Opin.Cell Biol. 8, 657-669
  - 54. Hughes, P.E., O'Toole, T.E., Ylanne, J., Shattil, S.J., and Ginsberg, M.H. (1995) J.Biol.Chem. 270, 12411-12417
- 55. Puzon McLaughlin, W., Yednock, T.A., and Takada, Y. (1996) J.Biol.Chem. 271, 16580-16585
  - 56. O'Toole, T.E., Katagiri, Y., Faull, R.J., Peter, K., Tamura, R., Quaranta, V., Loftus, J.C., Shattil, S.J., and Ginsberg, M.H. (1994) J.Cell Biol. 124, 1047-1059
- 25 57. De Melker, A.A., Kramer, D., Kuikman, I., and Sonnenberg, A. (1997) Biochem J 529-537
  - 58. Rojiani, M.V., Finlay, B.B., Gray, V., and Dedhar, S. (1991) Biochemistry 30, 9859-9866
- 59. Coppolino, M.G., Woodside, M.J., Demaurex, N.,
  30 Grinstein, S., St Arnaud, R., and Dedhar, S. (1997)
  Nature 386, 843-847
  - 60. Hynes, R.O. (1992) Curr.Opin.Genet.Dev. 2, 621-624
  - 61. Santoro, S.A. (1986) Cell 46, 913-920
- 62. Languino, L.R., Gehlsen, K.R., Wayner, E., Carter, W.G., Engvall, E., and Ruoslahti, E. (1989) J.Cell Biol. 109, 2455-2462

63. Yokosaki, Y., Monis, H., Chen, J., and Sheppard, D. (1996) J.Biol.Chem. 271, 24144-24150

## SEQUENCE LISTING

(1)	GENERAL	INFORMATION:

- (i) NUMBER OF SEQUENCES: 2
- (2) INFORMATION FOR SEQ ID NO. 1:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 3884 base pairs
    - (B) TYPE: nucleic acid and amino acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULAR TYPE: cDNA
  - (vi) ORIGINAL SOURCE:
    - (E) ORGANISM: human
    - (F) CELLTYPE: chondrocyte
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 1:

4	1				-+-			+				+			-+-			+			CCTG + GGAC	60
<b>a</b>									M	E	L	P	F	v	T.	н	L	F	L	P	L	-
	61				-+-			+				+			-+-			+			ATTC + TAAG	120
a				L					s						•					L	F	-
	121				-+-			+				+			-+-			+			ACAG	180
																	ACA.	ACC G	222 6	ACC G	TGTC	
а		P	G NTG	P GDT	P GCT	_	A ccc	_	_	G TTG	Y GGD	S TGG	ecc	ሁ ተጥር	Q AGG	H CGA	•		_	_	Q CGTT	-
	181				-+-			+		·		+			-+-			+			+	240
B		R	W	M	L	v	G	A	P	W	D	G	P,	s	G	D	R	R.	G	D <sub>.</sub>	v	-
	241		TATCGCTGCCCTGTAGGGGGGCCCACAATGCCCCATGTGCCAAGGGCCACTTAGGTGAC															300				
																•						
a		_	•	•	-		-	-	A TOTAL		•		P	_					L	_	D GTTA	-
	301				-+-			4				+			-+-			+			+	360
a		Y	Q	L	G	N	s	s	Н	P	A	v	N	M	н	L	G	M	s	L	L	-
	361																					
		GAGACAGATGGTGATGGGGGATTCATGGCCTGTGCCCCTCTCTGGTCTCGTGCTTGTGGC																				

ETDGDGGFMACAPLWSRACG -

		AG	CTC:	rgre	CTT	CAG'	TTC	TGG	GAT	ATG	TGC	CUG	161	GGA	160	TTU	AII			CA	t	480
	421	TC	GAG	ACA	GAA	GTC	AAG.	ACC	CTA	TÁC	acg	GGC	ACA	CCT	ACG	AAG'	TAA	GGT	CGG	AGT	CCT	400
a		s	S	v	F	s	s	G	I	С	A	R	V	D	A	5	F	Q	P	Q	G	-
	481																				GGAT	540
•	401																				CCTA	
a		s	L	A	P	T	A	Q	R	С	P	T	Y	M	D	V	V	I	V	L	D	-
	5 47	GG	CTC	CAA	CAG	CAT	CTA	ccc	CTG	GTC	TGA	AGT	TCA	GAC	CTT	CCT	ACG	AAG.	ACT	GGT	AGGG	600
	341																				rccc	
a		G	s	N	s	I	Y	P	W	s	E	v	Q	T	F	L	R	R	L	V	G	-
	601	AA	ACT	STT'	TAT'	TGA	ccc	AGA	ACA	GAT.	ACA	GGT	GGG	ACT	GGT	ACA	GTA	TGG	GGA	GAG	CCCT	660
	601																				GGGA	
a		ĸ	L	F	I	D	P	E	Q	I	Q	v	G	L	.v	Q	Y	G	E	s	P	-
	661	GT	ACA	rga:	GTG	GTC	CCT	GGG	AGA	TTT	CCG	AAC	GAA	.GGA	AGA	AGT	GGT	GAG	AGC	AGC	AAAG	720
	991	CA	TGT	ACT	CAC	CAG	GGA	ccc	TCT	AAA	GGC	TTG	CTT	CCI	TCT	TCA	CCA	CTC	TCG	TCG	TTTC	,,,
a		v	Н	E	W	S	r	G	D	F	R	T	ĸ	E	E	v	v	R	A	A	ĸ	-
	201	AA	CCT	CAG	TCG	GCG	GGA	GGG	ACG	AGA	AAC	AAA:	GAC	TGC	CCA	AGC	TAA	TAA	GGT	GGC	CTGC +	780
	121																				GACG	,00
a		N	L	s	R	R	E	G	R	E	T	ĸ	T	A	Q	A	I	M	v	A	С	-
	781	AC.	AGA.	AGG	GTT	CAG	TCA	GTC	CCA	TGG	GGG	CCG	ACC	CGA	GGC	TGC	ÇAG	GCT	ACT	GGT	GGTT	840
	701																				CCAA	
а		T	E	G	F	s	Q	s	H	·G	G	R	P	E	A	A	R	L	L	V	V	-
	0.41	GT	CAC	TGA	TGG	AGA	GTC	CCA	TGF	TGG	AGA	GGA	GCI	TCC	TGC	AGC	ACI	'AAA'	GGC	CTG	TGAG	900
	041	CA	GTG.	ACT.	ACC	TCT	CAG	GGI	ACI	ACC	TCI	CCI	'CGA	AGG	SACO	TCG	TGF	TTI	CCG	GAC	ACTC	
a		v	T	D	G	E	s	Н	D	G	E	E	L	P	A	A	L	Ķ	A	С	E	<b>-</b> ,
	901	GC	TGG.	AAG	AGT	GAC	ACG	CTA	TGG	GAT	TGC	AGI	CCI	TGC	STC	ACTA	CCI	CCC	GCG	GCA	GCGA	960
	901																				CGCT	
a		A	G	R	v	T	R	Y	G	I	A	· <b>v</b>	L	G	H	Y	L	R	R	Q	R	-
	061	GA	TCC	CAG	CTC	TTT	CCI	GAG	AG	\AA'	TAC	AAC	CAT	TG	CCAC	GTG!	ATC	CAGA	ATG!	GCG	SATTC	1020
	301	CT	AGG	GTC	GAG	AAA	GGZ	CTC	TC	TT?	AATO	TTC	ATA	AAC	GTO	CAC	'AGC	STCI	raci	rcgo	TAAG	
8		D	P	s	s	F	L	R	E	I	R	T	I	A	s	D	Þ	D	E	R	F	-
	100-	TT	CTT	CAA	TGT	CAC	AGA	ATG?	AGG(	CTG	TC	rga(	TG	ACA:	rtg'	rggi	\TG	CACT	rago	SAG	ATCGG	1080
	1021	AA	GAA	GTT	'ACA	GTG	TCI	'AC	CCC	GAC	GAG	ACT(	SACT	rgt	AAC	ACC:	rác	GTG	ATC	CTC	AGCC	
а		F	F	N	v	т	D	E	A	A	L	T	D	r	v	D	A	L	G	D	R	-

	1001		TTTGG																		1140
	1001		AAACC																		1140
a		1	F G	L	E	G	s	H	A	E	N	E	s	s	F	G	L	E	M	s	-
			ATTGG																		
	1141		TAACC																		1200
a		Ω	I G	F	s	T	н	R	L	ĸ	D	G	I	L	F	G	M	v	G	A	_
			GACTG																		
	1201		CTGAC																		1260
a		Y I	D W	G	G	s	v	L	W	Ĺ	E	G	G	H	R	L	F	P	P	R	_
		ATG	GCACT	'GGA	AGA	CGA	GTT(	CCC	CCC	rgcz	ACT(	3CA(	SAAG	CCA!	rgcz	AGC	CTAC	CTC	GG:	PTAC	
	1261		 CGTGA																		1320
a		M 2	A L	E	D	E	F.	P	P	A	L	Q	N	н	A	A	Y	L	G	Y	-
		TCT	GTTTC	TTC	CAT	GCT'	TTT	GCG	GGC.	rggz	ACG(	CCG	CTC	STT:	CTC	CTC	rgg	GC1	rcc:	rcga	
	1321		 DAAAG																		1380
a		s v	v s	S	M	L	L	R	G	G	R	R	L	F	L	s	G	A	P	R	
		TTT	AGACA	TCG	AGG	AAA	AGT	CAT	CGC(	CTTC	CCA	GCT:	AA1	GAAZ	AGA:	rgge	GGC:	rgto	EAG	GGTT	
	1381		 rctgt																		1440
a		F I	RH	R	G	ĸ	v	I	A	F	Q	L	ĸ	ĸ	D	G	A	v	R	v	_
		GCC	CAGAG	CCT	CCA	GGG	GGA	GCA(	GAT:	rgg:	rtci	ATA	CTT:	rgge	CAG:	rgae	GCT	CTGC	ccci	ATTG	
	1441		GTCTC																		1500
a		A (	g s	L	Q	G	E	Q	I	G	s	Y	F	G	s	E	L	С	P .	L	-
		GATA	ACAGA	TAG	GGA	TGG	AAC	AAC:	rga:	rgto	CTT	ACTI	rgto	GC.	rgco	CCC	CAT	STTO	CCT	GGGA	
	1501		 rgtct																		1560
a		D 1	r D	R	D	G	т	T	Đ	v	L	L.	v	A	A	P	M	F	L	G	_
	Çî.	ccc	CAGAA	CAA	GGA	AAC	AGG	ACG:	rgt	TA:	rgto	STA1	CTC	GTZ	AGG	CCA	GCA	GTC	CTT	GCTG	
	1561		 STCTT																		1620
а			O N																		_
			CTCCA																		
	1621			-+-			+							-+			+			+	1680
а			L Q																		_
•			CTCT																		
	1681			-+-			+-				<b></b> -			-+			+			+	1740
_			.unun															,		•	

	1741																				CCC	1800
	1,41																				GGG	
a		E	D	G	H	Q	G	A	L	Y	L	Y	H	G	T	Q	s	G	٧.	R	P	-
																					CGA	
	1801																		SAAA		GCT	1860
a		н	P						A		s		P	н	A	Ĺ	s	Y	F	G	R	_
_		-	- rgte		_		•						יבטי	rcro	GGT	CGA'	rgt	GGC	rgre	GGT	rGCC	
	1861				-+-			+				+			-+			+-				1920
									•													
a		S				-		÷		-	_		D			-					A	_
	1921	CAGGGGGCAGCCATCCTGCTCAGCTCCCGGCCCATTGTCCATCTGACCCCATCACTGGAG GTCCCCCGTCGGTAGGACGAGTCGAGGGCCGGGTAACAGGTAGACTGGGGTAGTGACCTC Q G A A I L L S S R P I V H L T P S L E GTGACCCCACAGGCCATCAGTGTGGTTCAGAGGGACTGTAGGCGGCGAGGCCAAGAAGCA CACTGGGGTGTCCGGTAGTCACACCAAGTCTCCCTGACATCCGCCGCTCCCGGTTCTTCGT V T P Q A I S V V Q R D C R R R G Q E A GTCTGTCTGACTGCAGCCCTTTGCTTCCAAGTGACCTCCCGTACTCCTGGTCGCTGGGAT															1980.					
		GT	ccc	CCG	TCG	GTA	GGA	CGA	GTC	GAG	GGC	CGG	GTA	ACA	GGT	AGA	CTG	GGG'	ragi	GAC	CTC	
a		Q	G,	A	A	I	L	L	s	S	R	P	1	V	H	L	T	P	S	L	E	-
	1981																					2040
	1301																					
a		V	T	P	Q	A	Ţ	s	v	v	Q	R	D	С	R	R	R	G	Q	E	A	<b>-</b> ,
	2041																				CTA	2100
a		v	С	L	T	A	A	L	С	F	Q	v	T	s	R	T	P	G	R	W	D	-
		CAC	CAZ	ATT(	CTA	CAT	GAG	GTT	CAC	CGC	ATC	ACT	GGA'	TGA	ATG	GAC	TGC	TGG	GGCI	ACG:	rgca	
	2101																				ACGT	2160
a		н	0										D		W	T	A	G	A	R.	A	_
		GCI	- ሚጥጥን	rga	TGG	CTC	TGG	CCA	GAG	GTT	STC	ccc	TCG	GAG	GCT	CCG	GCT	CAG	TGT(	GGG	GAAT	
	2161				-+-			+				+			-+-			+				2220
			-																v	G	N	-
<b>a</b>		A			G 			_					R.					•				
	2221				-+-			+				+			-+-			+				2280
																					ICAC	
а		V	T	С	E	Q	L	H	<b>F</b>	. <b>H</b>	V	L	D	T	S	D	Y	L	R	P	V	-
	2281																				TGAG	2340
																					ACTC	
a		A	L	T	v	T	F	A	L	D	N	T	T	ĸ	P	G	P	V	L	N	E	-
•																					CAAT	0400
	2341																				+ GTTA	2400
					_			_				_	_	_		_	_	_	_	_		

	2401																				GCC	2460
	2101																				CCGG	2400
a		E	С	v	T	D	L	v	L	Q	v	N	M	D	1	R	G	s	R	ĸ	A	_
		CC	ATT'	TGT	GGT	TCG.	AGG'	TGG	CCG	GCGC	GAAJ	GTG	сто	GTZ	ATC:	racz	ACT	CTC	GAG	AAC	CAGA	
	2461																				TCT	2520
a		P	P	v	V	R	G	G	R	R	ĸ	v	L	V	s	T	T	L	E	N	R	-
																					GCC	
	2521																				CCGG	2580
a		ĸ	E	N	A	Y	N	T	s	L	S	I	I	F	s	R	N	L	н	L	A	_
		AG	TCT	CAC!	rcc	TCA	GAG	AGA(	GAGO	CCC	ATA	AA.	GTC	GAZ	TGI	rgco	CGCC	ccc	TCI	GCI	CAT	
	2581																				GTA	2640
a		s	L	T	P	Q	R	E.	s	P	I	ĸ	v	E	С	A	A	P	s	A	н	_
		GC	CCG	GCT	CTG	CAG'	TGT	GGG	GCAT	rcci	rgto	TTC	CAC	ACT	rgg <i>i</i>	4GC0	CAAC	GTO	SACC	TTT	CTG	
	2641				•			•	CGTZ						•			•			+	2700
a									н													_
					•	•	•	٠												ACT	rGCC	
	2701				-+			+-			1				+			+-				2760
a																			L			_
_																						
	2761				-+			+-							-+			+-			+	2820
_																		_				
a			_	Ī.						•			_					_	_	_		
	2821	AGCAGTGACAGCCTGGAGAGAAATGGCACCCTTCAAGAAAACACAGCCCAGACCTCAGCC															2880					
a			_	_														•			Ε.	-
	2881				-+			+-			1				-+			+				2940
		CA	AGT	GG:	[AT]	ACC	CTG	GGA(	GGT	rca(	CCCF	\GG/	ACCG	GG'	rct"	PAA	GTT"	rtg	∃TG∤	\GA(	STCC	
a		V	H	P	Y	G	T	L	P	V	G	₽	G	P	E	F	K	T	T	L	R	-
	2941																				AGCT	3000
																					CGA	•
a		v	Q	N	L	G	С	Y	V	V	s	G	L	I	I	S	A	L	L	P	A	-
	3001																				AAGC	3060
	2001																				TTCG	
		v	ħ	ם	G	G	N	v	F	1.				^	v	т	T	N	N	Δ	•	_

	3061																				rcaa +	3120
																					AGTT	•
a		С	I	v	Q	N	L	T	E	P	P	G	P	P	v	H	P	E	E	L	Q	-
	2101																				rggg	2100
	3121																				ACCC	3180
a		н	T	N	R	L	N	G	s	N	T	Q	C	Q	v	v	R	С	H	L	G	-
	2101																				ATTT	2040
	3181																				AAAT	3240
а		Q	L	A	к	G	T	E	V	s	V	G	L	L	R	L	v	Н	N	E	F	-
	2044																				CGAA	2222
	3241																				GCTT	3300
a		F	R	R	A	ĸ	F	ĸ	s	L	T	v	v	s	, <b>T</b>	F	E	L	G	T	E	-
	3301																				GGTG	3360
	3301											•			-						CCAC	3300
a		E	G	s	V.	L	Q	L	. <b>T</b> .	<b>E</b>	Α.	s	R	W	s	·E	s	L	r	E	V	
	3361																				AGGG	3420
																					rccc	
a		V	Q	T	R	P	I	L	1	S	L	W	I	L	Í	G	s	v	L	. <b>G</b>	G	<b>-</b> .
·	3421																				TAAG	3480
		AA	CGA	GGA	CGA	ACG	AGA	GGA	ACA	GAA	GAC	GGA	CAC	CTT	CGA	ACC	GAA	GAA	ACG	GGT	ATTC	
a		L	L	L	L	A	L	L	<b>V</b> .	. <b>F</b>	С	L	W	K	L	G	F	F	A	H	K	-
	3481																				GGGT +	3540
		TT.	PTAC	GGG.	ACT	CCT	TCT	TTT	TTC	TCT	TCT	CTT	CAA	CCT	CGT'	TAC	TTA	CAT	CTT	ATT(	CCCA	
a	•						E								_			•				•
	3541																				GGGG +	3600
		GA'	CT	rtc.	AGG	AGG	GAC	CGT	CGA	AAG	AAG	TTC	TCT	GAA	CGT	ATT'	TTC	GTC	TCC	AAA	cccc	
																					TTGA	
	3601																				AACT	3660
																						•
	3661																				GAGC	3720
		GAZ	AAA	TC	AGG	ATC	CCT	ACG	ACG	ACC	TAR	CTC	TAC	TCC	GAA	ATG	GAG	TCT	GTT	CTT	CTCG	

3721	TGGCACCAAAACTAGCCATGCTCCCACCCTCTGCTTCCCTCCTCGTGATCCTGGTTC	379(
3121	ACCGTGGTTTTGATCGGTACGAGGGTGGGAGACGAGGAGGAGGAGGAGCACTAGGACCAAG	3700
3781	CATAGCCAACACTGGGGCTTTTGTTTGGGGTCCTTTTATCCCCAGGAATCAATAATTTTT+ GTATCGGTTGTGACCCCGAAAACAAACCCCAGGAAAATAGGGGTCCTTAGTTATTAAAAA	3840
3841	TTGCCTAGGAAAAAAAAAGCGGCCGCGAATTCGATATCAAGCT	

(2)																						
	<b>i</b> )	L)	S	EQ	UEI	NCE	C	HA	RA	CTE	ERI	ST	IC	S:								
				(A)	]	LEN	IGI	Ή:	3	779	b	as	e p	bai	rs							
				(B)		ryf	E:	n	uc	lei	.d	ac.	id	an	d	am:	inc	a	ci	d		
						STF																
						rof											•			,		
				(E)			-															
	/ 4	Ĺ)		-		FTT 7	ND.	πv	DE	. ,	יארו -	τ 20.					•					
	•	•										ın					٠					
·	.13	/i)																		*		
						OR									_							
				(B)		CE	ىلىل	Ί. Χ .	PE:	C	no	naı	coc	:yt	е							
			_										_		* -		_	•				
	( )	ki)	S	EQ	UE:	NCE	SI	DES	CR	1 P'.	LTC	)N:	S.	ĿQ	ID	N	υ.	2:	;			
									•						•							
		ሮልር	e CT(	CNG	מממ	cce	<b>ስ</b> ጥር	A CC	ፖልጥ	GGA	АСТ	CCC	CTT	CGT	CAC'	rca(	CCT	STT	CTT	300	CTG	
	1				-+-			+				+			-+			+			+	60
		GTO	CAC	GTC'	TTT	GGC'	TAG	TCC	GTA	CCT	TGA	GGG	GAA	GCA	GTG	AGT	GGA	CAA	GAA	CGG	GGAC	
									M	F	τ.	ъ	E-	17	· •	u	<b>T</b> .	F	7.	D	L	_
									. 144	-		•	Ŀ	•	•	••	_	٠	_	•	_	
		GTO	STT(	CCT	GAC	AGG'	TCT	CTG	CTC	CCC	CTT	TAA	CCT	GGA'	TGA	ACA	TCA	CCC	ACG	CCT	ATTC	
	61				-+-			+				7 TT			~+~·	 rcr	и С.Д. 	+ ccc			+ TAAG	120
		CAC	JAA	<b>GUA</b>	CIG	100.	АСН	GAC	GAG	999	GAA	WII	GGA	CC1.	HCI		ng I	<b>66</b> 6	160	JOA	IAAG	
а		v	F	L	T	·G	L	С	S	P	F	N	L	D	E	H	H	P	R	L	F	· <b>-</b>
		~~1				202	<b>.</b>	mc3	3 mm	mcc	አ መ አ	C) C	ጥረጥ	ርጥጥ	አሮአ	ארא	ጥረጥ	ጥርር	ccc	TGG:	ACAG	•
	121	ÇC	AGG(	GCC.	ACC	AGA 	AGC 	TGA		166			1G1	<del></del>	-+-			+		100	+	180
•		GG?	rcc	CGG	TGG	TCT	TCG	ACT	'TAP	ACC	TAT	GTC	ACA	GAA	TGT	TGT	ACA	ACC	ccc	ACC'	TGTC	
42		_	_		ъ		7	r	F	c	v	e	3.7	τ.		น	v	G	G	· .	Q	_
a.		2	G	P	P	£	A	E	£	: 6	•	3	•	_	¥	••	•	•	Ŭ		¥	
		CGZ	ATG	GAT	GCT	GGT	GGG	CGC	ccc	CTG	GGA	TGG	GCC	TTC	AGG	CGA	CCG	GAG	GGG	GGA	CGTT	
	181				-+-			+		CAC		+	ccc		-+- TCC	 CCT	ccc 	+ ርጥር		 ССТ	+ GCAA	240
a		R	W	M	L	v	G	A	P	W	D	G	P	S	G	D	R	R	G	D	V	-
		m h r	PCC.	OT (*)		ጥርጥ	D.C.C	cco	ccc	ירר <i>י</i>	ממי	тсс	יררר	אייני	ጥርር	مم	GGG	CCA	СТТ	AGG	TGAC	
	241	1A.			-+-							+			-+-			+			+	300
		AT	AGC	GAC	GGG	ACA	TCC	ccc	CCC	GGT	GTT	'ACG	GGG	TAC	ACG	GTT	CCC	GGT	GAA	TCC	ACTG	
		•	_	_	_	٠,,	_	_			M		ъ	_	7.	v	c	u	·T.	c	ח	
a		Y	R.	C	Ρ.	V	G	G	A	н	14	A	. ₽	C	A	r	Ģ	п	L	G	ט	_
		TAC	CCA	ACT	GGG	AAA	TTC	TA	TC	TCC	TGC	TGT	GAA	TAT	GCA	CCT	GGG	GAT	GTC	TCT	GTTA	
	301				-+-							+			-+-			+			+	360
•		TA	3GT	TGA	CCC	TTT	AAC	TAG	AGI	AGG	ACG	ACA	CTI	WIH	CGI	GGH		CIA	CAC	MOM	CAAT	
a		Y	Q	L.	G	N	s	s	н	₽	A	v	N	M	H	L	G	M	S	L	L	-
		·																			mccc	
	361	GA(	GAC.	AGA 	TGG	TGA	TGO	GGG	AT	CAT	'GGC	CTG	TGC	CCC	TCT	CTG	GIC	TCG	TGC	116	TGGC	420
	201	CT	CTG	TCT	ACC	ACT	ACC	CCC	TAI	GTA	CCG	GAC	ACG	GGG	Aga	GAC	CAG	AGC	ACG	AAC	ACCG	
•																						
a		E	T	D	G	D	G	G	F	M	A	С	A	P	L	W	S	R	A	С	G	-
		AG	СТС	TGT	CTT	CAG	TTC	TG	GAT	OTA1	TGC	ccc	TGI	'GGZ	TGC	TTC	:ATT	CCF	GCC	TCA	GGGA	
	421				-+-							-+			-+-						+	480
		TC	GAG	ACA	GAA	GTC	:AAC	SAC	CT	ATAC	CACC	GGG	CAC	CCI	ACG	AAC	'AT	GGI	CGG	AGT	CCCT	

a ·		S	S	V	F	S	S	G	I	С	A	R	V	D	A	S	F	Q	P	Q	G	-
	481																					540
	401																		ACA	SAA(	CCTA	
a		s	L	·A	P	T	A	Q.	R	С	P	T	Y	M	D	<b>v</b> .	v	I	<b>v</b> .	L	D.	<b>-</b>
	5.43	GĢ	CCTGGCACCCACTGCCCAACGCTGCCCAACATACATGGATGTTGTCATTGTCTTGAY  GGACCGTGGGTGACGGGTTGCGACGGGTTGTATGTACCTACAACAGTAACAGAACCTY  L A P T A Q R C P T Y M D V V I V L D  CTCCAACAGCATCTACCCCTGGTCTGAAGTTCAGACCTTCCTACGAAGACTGGTAGG  GAGGTTGTCGTAGATGGGGACCAGACTTCAAGTCTGGAAGGATGCTTCTGACCACTCCC  S N S I Y P W S E V Q T F L R R L V G  ACTGTTTATTGACCCAGAACAGATACAGGTGGGACCACTGCAAGACTGGTAGGGAAGACTGGTATGGGGAAGACCCTTCAGACAATAACTGGGTCTTGTCTATGTCCACCCTGACCATGCATACAGTATGGGGAAGACCCTTCAGACAAATAACTGGGTCTTGTCTATGTCCACCCTGACCATGCATACCCCTCTCGGGG  L F I D P E Q I Q V G L V Q Y G E S P  ACATGAGTGGTCCCTGGGAGATTTCCGAACGAAGGAAGAAGTGGTGAGAGCAGCAAAA  TGTACTCACCAGGGACCCTCTAAAGGCTTGCTTCCTTCTTCACCACTCTCGTCGTTT  H E W S L G D F R T K E E V V R A A K  CCTCAGTCGGCGGGAGGGACGAGAAACAAAGACTGCCCCAAGCAATAATGGTGGCCTG  GGAGTCAGCCGCCCTCCCTGCTCTTTGTTTCTGACGGGTTCGTTATTACCACCGGAC  L S R R E G R E T K T A Q A I M V A C  CAGAGAGGGTTCAGTCAGTCCCATGGGGGCCCGACCGAGGCTACTGGTGGT  TCTTCCCAAGTCAGTCAGGTACCCCCGGCTGCCCAAGGCTACTGGTGGT  TCTTCCCCAAGTCAGTCAGGGTACCCCCGGCTCCCGACGGTCCGAGGCTACTGGTGGT  TCTTCCCCAAGTCAGTCAGGGTACCCCCGGCTCCCGACGGTCCCGAGGCTACTGGTGGT  CGTGACTACCTCTCAGGGTACCCCCGGCTCCCGAGGACCTAAAAGGCCTGTGA  GGGACGTATGGAGAACAACAAGACAAG													AGGG	600					
	241																					
a		G	S	N	S	I	Y	P	W	S	E	V	Q	T	F	L	R	R	L	V	G	-
	601																					660
	001																					
a		ĸ	L	F	I	D	P	E	Q	I.	Q	V	G	L	V	Q	Y	G	E	S	P	<b>-</b>
	661	GT	ACA	TGA	GTG	GTC	CCT	GGG	AGA	TTT	CCG	AAC	GAA	GGA.	AGA	AGT	GGT	GAG	AGC	AGC	AAAG	720
	001	CA'	TGT	ACT	CAC	CAG	GGA	ccc	TCT	'AAA	GGC	TTG	CTT	CCT	TCT	TCA	CCA	CTC	TCG'	ŢCG'	TTTC	, 20
a		v	н	E	W	s	L	G	D	F	R	T	ĸ	E	E	v	V	R	A	A	ĸ	-
	701	ÄA	CCT	CAG	TCG	GCG	GGA	GGG	ACG	AGA	AAC	AAA	GAĊ	TGC	CCA	AGC	AĄT	AAT	GGT	GGC	CTGC	780
	721																					
a		N	L	s	R	Ŕ	E	G	R	E	T	K	T	A	Q	A	I	M	V	A	С	<del>-</del>
	701	AC	AGA	AGG	GTT	CAG	TCA	GTC	CCA	TGG	GGG	CCG	ACC	CGA	GGC	TGC	CAG	GCT	ACT	GGT	GGTT	840
•	101																					
a		T.	E	G	F	s	Q	s	H	G	G	R	P	E	A	A	R	L	L	v	v	-
	841		CAC	TGA	TGG	AGA	GTC	CCA	TGA	TGG	AGA	.GGA	GCT	TCC	TGC	AGC	ACI	'AAA'	GGC	CTG	TGAG	900
	011																					
a		v	T	D	G	E	S	H	D	G	. <b>E</b>	E	L	P	A	A	L	K	A	С	E	-
	901																					
	,,,,																					
a		A	G	R	V.	T	R	, <b>Y</b>	G	I	A	V	L	G	H	Y	L	R	R	Q	·R	-
	961																					
	, , ,																					
a		D	P	S	s	F.	L	R	E	I	R	T	I	A	s	D	P	D	E	R	F	-
	1021	TT	ACAGAAGGGTTCAGTCAGTCCCATGGGGGCCCGACCCGA														TCGG	1080				
a		F	F	N	V	T	D	E	A	A	L	T	D	I	v	D	A	L	G	D	R	-
	1003																					
	1091																				CAGA	

a		I	F	G	L	E	G	s	Н	A	E	N	E	s	s	F	G	L	E	M	S	-
	1141																					1200
	1141																					
a		Q	1	G	F	S	T	H	R	L	K	D	G	I	L	F	G.	M	<b>V</b>	G	A	-
	1201		CAGATTGGTTTCTCCACTCATCGGCTAAAGGATGGGATTCTTTTTGGGATGGTGGGGGCCCCTTAACCAAAGAGGTGAGTAACCCAACCCCCGG GTCTAACCAAAGAGGTGAGTAGCCGATTTCCTACCCTAAGAAAAACCCTACCACCCCCGG Q I G F S T H R L K D G I L F G M V G A TATGACTGGGGAGGCTCTGTGCTATGGCTTGAAGGAGGGCCACCGCCTTTTCCCCCCACGA ATACTGACCCCTCCGAGACACGATACCGAACTTCCTCCGGTGGCGGAAAAGGGGGGTGCT Y D W G G S V L W L E G G H R L F P P R ATGGCACTGGAAGACGAGTTCCCCCCTGCACTGCAGAACCATGCAGCCTACCTGGGTTAC TACCGTGACCTTCTGCTCAAGGGGGGACGTGACGT															1260				
а		Y	D	W	G	G	Ś	V	L	W	L	E	G	G	H	R	L	F	P	₽	R	-
	1261																					1320
		TA	CCG:	rga	CCT'	TCT	GCT	CAA	GĠG	GGG	ACG:	rga(	CGTC	TTC	GT?	ACGI	CGC	TAE	GAC	CCCI	ATG	
a		M	A	L	E	D	E	F	P	P	A	L	Q	N	H	A	A	Y	L	G	Y	-
																						1380
																				AGG	AGCT	
a																						-
	1381				-+-			+				+			-+			+-			+	1440
а										٠	٠	_								-		•
	1441				-+			+				+			-+			+-			+	1500
a																						_
•						-																
	1501				-+-			+				+			-+			+-			+	1560
a		D	T	D	R	D	G	T	T	D	v	L	L.	v	A	A	P	M	F	L	G	_
		CC	CCAC	SAAC	CAA	GGA	AAC	AGG	ACG'	TGT'	TTA!	rgte	rate	CTC	GT#	AGG	CAC	GCA	STC	CTT	CTG	
	1561																		CAG	GAAG		1620
a		P	Q	N	ĸ	E	T	G	R	V	. <b>Y</b>	v	<b>Y</b> .	L	v	G	Q	Q	s	L	L	-
	1.601																					1 600
	1621																					1680
a		T	L	Q	G	T	L	Q	P	E	P	P	Q	D	A	R	F	G	F	A	M	-
	1601																					1740
	1001		ARATCTGTAGCTCCTTTTCAGTAGCGGAAGGTCGAATTCTTTCT															1740				
a		G	A	L	P	D	L	Ŋ	Q	D	G	F	A	D	v	A	V	G.	A	P	L	-
	1741																					1800

a		E	D	G	Н	Q	G	A	L	Y	L	Y	H	G	T	Q	S	G	V	R	P	-
	1801				-+-			+				+			-+-			+			+	1860
		GT.	AGG	ACG	GGT	CTC	CTA	ACG	ACG	ACG	GAG	GTA	CGG'	rgt.	ACG	GGA	GTC	GAT	GAA	ACC	GGCT	
a		H	P	A	Q	R	I	A	A	A	S <sub>.</sub>	M	P .	H	A	L	S	Y	F	G	R	-
	1061		ATCCTGCCCAGAGGATTGCTGCTGCCTCCATGCCACATGCCCTCAGCTACTTTGGCCG  PAQRIAAAS MPHALSYFGR  CACACCTACCAGCCGATCTAGACCTACCTCTACTAGACCAGCTACACCGACCACCG  CACACCTACCAGCCGATCTAGACCTACCTCTACTAGACCAGCTACACCGACACCCACG  VDGRLDLDGDDLVDVAVGA  CACACCTACCAGCCGATCTGCATCGAGGGCCCATTGTCCATCGACCCCACCGACCCCACGGCCCAGCCCACCGTCCCGGTAGGACGAGCCCACCGGCCCATCGTCCGGTAGGACGAGCCACCCAC														1020					
	1001																					1920
a		s	v	D	G	R	Ŀ	D	L	D	G	D	D	L	v	D	v	A	v	G	A	<b>-</b> .
		CA	GGG	GGC	AGC	CAT	CCT	GCT	CAG	CTC	CCG	GCC	CAT!	rgte	CCA:	гст	GAC	ccc	ATC.	ACT	GGAG	
	1921																					1980
a		0	G	A	A	ī	I.	L	s	s	R	Þ	T	v	Ħ	т.	T	p	s	t.	£	_
_												•						_			_	
	1981				-+-			+				+			-+-			+			+	2040
•				-					ACA		•								GGT		TCGT	
a		V	T	P	Q	A	I	S	V	V	Q	R	D.	C	R	R	R	G	Q	E	A	-
	2041																					2100
		CA	GAC	AGA	CTG	ACG'	TCG	GGA	AAC	GĄA	GGT'	rca(	CTG	GAG	GGC	ATG	AGG	ACC.	AGC	GAC	CCTA	
a		v	С	L	T	A	A	L	С	. <b>F</b>	Q	٧.	T	S	R	T	P	G	R	Ŵ	D	-
	2101				CTA	CAT	GAG	GTT	CAC	CGC	ATC	ACT	GA.	rga.	ATG	GAC:	rgc	TGG	GGC.	ACG	TGCA	2160
	2101																					2100
a		н	Q	F	Y	M	R	F	T	A	s	L	D	E	W	T	A	G	A	R	A	-
																			TGT	GGG		
	2161																		ACA	CCC		2220
a		A	F	D	G	s	G	Q	R	L	s	P	R	R	L	R	L	s	v	G	N	_
		GT	CAC:	rtg:	rga(	GCA	GCT	ACA	CTT	CCA:	rgto	CTC	GA1	raci	ATC	AGA'	AT1	CCT	CCG	GCC	AGTG	
	2221																					2280
A																						_
٠.																						_
	2281				-+			+				+			-+			+			+	2340
															•							
a		A	L	T	V	T	F	A	L	. D	N	T	T	K	P	G	P	V	L	N	E	-
	2341																				CAAT	2400
																					GTTA	
a		G	S	P	T	s	Ĭ.	Q	ĸ	L	v	P	F	s	ĸ	D	С	G	P	D	N	-
																					GCC	
	2401																				ccee	

a		E	C V	T	D	L	V	L	Q	V	N	M	D	1	R	G	S	R	K	A	-
	2461			+-			+				+			-+			+				2520
_			AAAC F V												T.	TTG.	AGA:	E		R	
а	2521	AAG	GAAA	ATGC	ATT:	CAA	TAC	GAG	CCT	GAG	TAT +	CAT	CTT	CTC:	TAG	AAA	CCT	CCA	CCT	GGCC	2580
			CTTT																		
а			E N																		-
	2581			+-			+				+			-+-			+			AGTA	2640
a		s	L I	P	Q	R	E	s	P	Ĭ	ĸ	v	E	Ċ	A	A	P	s	,A	н	-
	2641		CGGC																		2,700
		CGG	GCCG	AGAC	GTC	ACA	CCC	CGT	AGG	ACA								CTG	GAA	AGAC	_,, 00
а			R I	-			G	H	_	V 	_	Q						T	F	_	-
	2701	CTAGAGTTTGAGTTTAGCTGCTCCTCTCTCCTGAGCCAGGTCTTTGGGAAGCTGACTGCC  O1+										2760									
a			E I																		-
	2761	AGC	AGTO	ACAC	CCI	GGA	GAG	AAA	TGG	CAC	CCT	TCA	AGA.	AAA -+-	CAC	AGC	CCA	GAC	СТС	AGCC	2820
		TCG	TCAC	TGT	CGGA	CCT	CTC	TTT	ACC	GTG ·	GGA	AGT	TCT'	TTT	GTG	TCG	GGT	CTG	GAG	TCGG	
a			s I	_								Q					Q CCN	T	S	A TCDC	-
	2821			+-							+			-+-			+			TGAG + ACTC	2880
a		Y	1 (	Y	E	P	н	L	L	F	s	s	E	s	T	ŗ	н	R	Y	E	-
	2881			+-							+			-+-			+			CAGG + .GTCC	2940
a			н 1																		-
	2941			+-							+			-+-			+			GCAT	3000
a		T	n i	ı A	s	С	I	v	Q	N	L	T	E	P	P	G	P	P	v	н	-
	3001			+-							+			-+-			+				3060
								• •	• •	• •										v V	
а		AGG	TGC	CACC	rtge	GC#	AGCI	r GGC	CAAZ	AGGG	GAC	TGA	GGT	CTC	TGT	TGG	ACI	rran	GAG	GCTG	
	3061	AGGTGCCACCTTGGGCAGCTGGCAAAGGGGACTGAGGTCTCTGTTGGACTATTGAGGCTG+ 31											3120								

a		R	С	H	L	G	Q	L	A	ĸ	G	T	E	V	S	V	G	L	L	R	L	-									
	3121	GTTCACAATGAATTTTTCCGAAGAGCCAAGTTCAAGTCCCTGACGGTGGTCAGCACCTTT															3180														
		CA	AGT	STGTTACTTAAAAAAGGCTTCTCGGTTCAAGTTCAGGGACTGCCACCAGTCGTGGAAA																											
a		V	H <sub>.</sub>	N	E.	F	F	R	R	A	K	F	K	S	L	T	V	V	S	T	F	-									
•	3181		GAGCTGGGAACCGAAGAGGGCAGTGTCCTACAGCTGACTGA															3240													
а		E		G								-	L									- '									
	3241	AGCCTCTTGGAGGTGGTTCAGACCCGGCCTATCCTCATCTCCCTGTGGATCCTCATAGGC														3300															
		TCGGAGAACCTCCACCAAGTCTGGGCCGGATAGGAGTAGAGGGACACCTAGGAGTATCCG																													
a		s	L	L	E	V	V	Q	T	R	P	Ţ	L	I	S	L	W	I	L	I	G	-									
	3301																				TGGC	3360									
	. 3301																				ACCG	2200									
a		s	v	L	G	G	L	L	r	L	À	L	L	v	F	С	L	W	K	L	G	-									
	3361	TTCTTTGCCCATAAGAAAATCCCTGAGGAAGAAAAAAGAGAAGAGAAGATTGGAGCAATGA												2420																	
																					TACT	3420									
a		F	·F	A	н	ĸ	K	I	P	E	E	E	ĸ	R	E	E.	ĸ	L	E	Ò											
	3421																				AAATA										
																					ATTT	+ 3480 T									
		3.0	a	<b>3.CC</b>	mmer				CDT	ccc	יאריא	nc.	אמכר	rcc	·CTC	ጥርር	יים מי	ימירי	ייירר	יררז	አር <u>አ</u> ርር	:ACC									
	3481	AGCAGAGGTTTGGGGGCTCAGATGGGACAAGAAGCCGCCTCTGGACTATCTCCCCAGACC  TCGTCTCCAAACCCCCGAGTCTACCCTGTTCTTCGGCGGAGACCTGATAGAGGGGTCTGG											3540																		
		TC	GTC	TCC	AAA	CCC	CCC	AGT	CTA	ıccc	rGi	TCI	TCG	الحال	GAG	MCC	TGF	MC	MGG	1001	CIGG	iG									
			CAG	ССТ																	TACC										
	3541		GTC	GGA	•			-													ATGG	3600									
		ጥሮ	DCD.	<b>~</b> AA	CD.	GAG	: ርጥር	cci	 VCCE		ג ח <i>יי</i>	vec (	ን ጥ ሬ	CTC	CCZ	CCC	CCCTCTGCTTCCCTCCTCC														
	3601				-+-			1				+			-+-						+ GGAGG	3660									
		AG	TCI	GII	CII	CIC	,GA(	.061	GGI		CAI	CGC	JIAC	.GAC		. 660	ngr	ıcar	2100	JOAL	Janoo										
	2661	TCGTGATCCTGGTTCCATAGCCAACACTGGGGCTTTTGTTTG																													
	3661																				GGTC										
		GA	ATC	TAA:	'AA'	TTT	TTT	rgco	CTAC	GAZ	<b>LAA</b>	LAA.	AA.	SCGG	ccc	SCG7	\AT	rcg <i>i</i>	ATA:	rcaj	AGCT										
	3721											-+			+-							3779									

- (2) INFORMATION FOR SEQ ID NO. 3:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 143 base pairs
    - (B) TYPE: nucleic acid and amino acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (iii) MOLECULAR TYPE: cDNA
  - (vi) ORIGINAL SOURCE:
    - (A) ORGANISM: human
    - (B) CELLTYPE: chondrocyte
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 3:

NdeI

GGGGCATATGGTTCAGAACCTGGGTTGCTACGTTGTTTCCGGTCTGATCATCTCCGCTCT

1 -----+ 60
CCCCGTATACCAAGTCTTGGACCCAACGATGCAACAAAGGCCAGACTAGTAGAGGCGAGA

b GHMVQNLGCYVVSGLJISAL-

GCTGCCGGCTGTTGCTCACGGTGGTAACTACTTCCTAAGCTTGTCCCAGGTTATCAGCGG
61 -----+ 120
CGACGGCCGACAACGAGTGCCACCATTGATGAAGGATTCGAACAGGGTCCAATAGTCGCC

b LPAVAHGGNYFLSLSQVISG-

BamHI

b L V P R G S P \_-

15

25

30

## CLAIMS

- 1. A recombinant or isolated integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity.
- 2. A process of producing a recombinant integrin subunit all comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or homologues or fragments thereof having similar biological activity, which process comprises the steps of
- a) isolating a polynucleotide comprising a nucleotide sequence coding for an integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity,
- b) constructing an expression vector comprising the isolated polynucleotide,
- c) transforming a host cell with said expression
   vector,
  - d) culturing said transformed host cell in a culture medium under conditions suitable for expression of integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, in said transformed host cell, and, optionally,
  - e) isolating the integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, from said transformed host cell or said culture medium.
  - 3. A process of providing an integrin subunit  $\alpha 10$ , or homologues or fragments thereof having similar biological activity, whereby said subunit is isolated from a cell in which it is naturally present.
- An isolated polynucleotide comprising a nucleotide coding for an integrin subunit α10, or for homologues or fragments thereof, which polynucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or suitable parts thereof.

5. An isolated polynucleotide or oligonucleotide which hybridises to a DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit  $\alpha 1$ .

5

10

15

30

35

- 6. A vector comprising a polynucleotide or oligonucleotide coding for an integrin subunit α10, or homologues or fragments thereof, which polynucleotide or oligonucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof.
- 7. A vector comprising a polynucleotide or oligonucleotide which hybridises to a DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit  $\alpha 1$ .
- 8. A cell containing the vector as defined in any one of claims 6 and 7.
- 9. A cell generated by the process in claim 2, in which a polynucleotide or oligonucleotide coding for an integrin subunit α10, or homologues or fragments thereof, which polynucleotide or oligonucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof has been stably integrated in the cell genome.
  - 10. Binding entities having the capability of binding specifically to integrin subunit  $\alpha 10$  comprising the amino acid sequence of SEQ ID No. 1 or SEQ ID No. 2, or to homologues or fragments thereof.
  - 11. Binding entities according to claim 10, which are chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.
  - 12. A recombinant or isolated integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the

5

10

15

20

35

subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, and homologues and fragments thereof having similar biological activity.

- 13. A recombinant or isolated integrin heterodimer according to claim 12, wherein the subunit  $\beta$  is  $\beta 1.$
- 14. A process of producing a recombinant integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , in which the subunit  $\alpha 10$  comprises the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, and homologues and fragments thereof, which process comprises the steps of
- a) isolating one polynucleotide comprising a nucleotide sequence coding for a subunit  $\alpha 10$  of an integrin heterodimer and, optionally, another polynucleotide comprising a nucleotide sequence coding for a subunit  $\beta$  of an integrin heterodimer, or polynucleotides or oligonucleotides coding for homologues or fragments thereof having similar biological activity,
- b) constructing an expression vector comprising said isolated polynucleotide coding for said subunit  $\alpha 10$  optionally in combination with an expression vector comprising said isolated nucleotide coding for said subunit  $\beta$ ,
- c) transforming a host cell with said expression vector or vectors,
- d) culturing said transformed host cell in a culture medium under conditions suitable for expression of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof having similar biological activity, in said transformed host cell, and, optionally,
  - e) isolating the integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or homologues or fragments thereof having similar biological activity, or the  $\alpha 10$  subunit thereof from said transformed host cell or said culture medium.
  - 15. A process of providing an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta\text{,}$  or homologues

10

15

or fragments thereof having similar biological activity, whereby said integrin heterodimer is isolated from a cell in which it is naturally present.

- 16. A cell containing a first vector, said first vector comprising a polynucleotide or oligonucleotide coding for a subunit  $\alpha 10$  of an integrin heterodimer, or for homologues or parts thereof having similar biological activity, which polynucleotide or oligonucleotide comprises the nucleotide sequence shown in SEQ ID No. 1 or SEQ ID No. 2 or parts thereof, and a second vector, said second vector comprising a polynucleotide or oligonucleotide coding for a subunit  $\beta$  of an integrin heterodimer, or for homologues or fragments thereof.
- 17. Binding entities having the capability of binding specifically to the integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof, or a subunit  $\alpha 10$  thereof, having similar biological activity.
- 18. Binding entities according to claim 17, wherein 20 the subunit  $\beta$  is  $\beta 1.$ 
  - 19. Binding entities according to claim 17 or 18, which are chosen among the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, and fragments thereof.
- 20. A fragment of the integrin subunit α10, which fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 21. A fragment according to claim 20, which is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
  - 22. A fragment according to claim 20, which comprises the amino acid sequence from about amino acid No. 952 to about amino acid no. 986 of SEQ ID No. 1.
- 23. A fragment according to claim 20, which is a peptide comprising the amino acid sequence from about

15

20

25

35

amino acid No. 140 to about amino acid no. 337 of SEQ ID No. 1.

- 24. A method of producing a fragment of the integrin subunit  $\alpha 10$  as defined in any one of claims 20-23, which method comprises a sequential addition of amino acids containing protective groups.
- 25. A polynucleotide or oligonucleotide coding for a fragment of the integrin subunit  $\alpha 10$  as defined in any one of claims 20-23.
- 26. Binding entities having the capability of binding specifically to a fragment of the human integrin subunit  $\alpha 10$  as defined in any one of claims 20-23.
  - 27. Binding entities according to claim 26, which are chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, and fragments thereof.
  - 28. A process of using an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or a homologue or fragment of said integrin or subunit having similar biological activity, as a marker or target molecule of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.
  - 29. A process according to claim 28, whereby said fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 30. A process according to claim 29, whereby said fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
  - 31. A process according to claim 29, whereby said fragment comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEQ ID No. 1.
  - 32. A process according to claim 29, whereby said fragment comprises the amino acid sequence from about

WO 99/51639 PCT/SE99/00544

amino acid no. 140 to about amino acid no. 337 of SEQ ID No. 1.

5

25

35

- 33. A process according to claim 28, whereby the subunit  $\beta$  is  $\beta$ 1.
- 34. A process according to claim 28, whereby said cells are chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.
- 35. A process according to any one of claims 28-34, which process is used during pathological conditions involving said subunit  $\alpha 10$ .
  - 36. A process according to claim 35, which pathological conditions comprise damage of cartilage.
- 37. A process according to claim 36, which patho15 logical conditions comprise trauma, rheumatoid arthritis
  and osteoarthritis.
  - 38. A process according to any one of claims 28-34, which is a process for detecting the formation of cartilage during embryonal development.
- 39. A process according to any one of claims 28-34, which is a process for detecting physiological or therapeutic reparation of cartilage.
  - 40. A process according to any one of claims 28-34, which is a process for selection and analysis, or for sorting, isolating or purification of chondrocytes.
  - 41. A process according to any one of claims 28-34, which is a process for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes.
- 30 42. A process according to any one of claims 28-34, which is a process for in vitro studies of differentiation of chondrocytes.
  - 43. A process of using binding entities having the capability of binding specifically to an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homo-

15

20

25

30

35

logues or fragments thereof having similar biological activity, as markers or target molecules of cells or tissues expressing said integrin subunit  $\alpha 10$ , which cells or tissues are of animal including human origin.

- 44. A process according to claim 43, whereby said fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 45. A process according to claim 43, whereby said
  10 fragment is a peptide comprising the amino acid sequence
  KLGFFAHKKIPEEEKREEKLEQ.
  - 46. A process according to claim 43, whereby said fragment comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEO ID No. 1.
  - 47. A process according to claim 43, whereby said fragment comprises the amino acid sequence from about amino acid no. 140 to about amino acid No. 337 of SEQ ID No. 1.
  - 48. A process according to claim 43, whereby the subunit  $\beta$  is  $\beta1$ .
    - 49. A process according to any one of claims 43-48, which is a process for detecting the presence of an integrin subunit  $\alpha 10$  comprising the amino acid sequence shown in SEQ ID No. 1 or SEQ ID No. 2, or of an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or of homologues or fragments thereof having similar biological activity.
    - 50. A process according to any one of claims 43-48, which process is a process for determining the differentiation-state of cells during embryonic development, angiogenesis, or development of cancer.
    - 51. A process for detecting the presence of an integrin subunit  $\alpha 10$ , or of a homologue or fragment of said integrin subunit having similar biological activity, on cells, whereby a polynucleotide or oligonucleotide chosen from the group comprising a polynucleotide or oligo-

15

25

nucleotide shown in SEQ ID No. 1 is used as a marker under hybridisation conditions wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit  $\alpha$ 1.

- 52. A process according to claim 51, whereby said cells are chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.
- 53. A process according to claim 51, whereby said

  10 fragment is a peptide chosen from the group comprising
  peptides of the cytoplasmic domain, the I-domain and the
  spliced domain.
  - 54. A process according to claim 53, whereby said fragment is a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
  - 55. A process according to claim 53, whereby said fragment comprises the amino acid sequence from about amino acid No. 952 to about amino acid no. 986 of SEQ ID No. 1.
- 56. A process according to claim 53, whereby said fragment comprises the amino acid sequence from about amino acid No. 140 to about amino acid No. 337 of SEQ ID No. 1.
  - 57. A process according to any one of claims 43-48, which is a process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration or in therapeutic and physiological reparation of cartilage.
- 58. A process according to claim 57, wherein the pathological conditions are any pathological conditions involving the integrin subunit  $\alpha 10$ .
  - 59. A process according to claim 58, whereby said pathological conditions are rheumatoid arthritis, osteoarthrosis or cancer.
- 35 60. A process according to claim 57, whereby said cells are chosen from the group comprising chondrocytes,

10

20

smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.

- 61. A process for determining the differentiation-state of cells during development, in pathological conditions, in tissue regeneration and in therapeutic and physiological reparation of cartilage, whereby a polynucleotide or oligonucleotide chosen from the nucleotide sequence shown in SEQ ID No. 1 is used as a marker under hybridisation conditions wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit  $\alpha$ 1.
- 62. A process according to claim 61, whereby said polynucleotide or oligonucleotide is a polynucleotide or oligonucleotide coding for a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the spliced domain.
- 63. A process according to claim 62, whereby said polynucleotide or oligonucleotide is a polynucleotide or oligonucleotide coding for a peptide comprising the amino acid sequence KLGFFAHKKIPEEEKREEKLEQ.
- 64. A process according to claim 62, whereby said peptide comprises the amino acid sequence from about amino acid no. 952 to about amino acid no. 986 of SEQ ID No. 1.
- 25 65. A process according to claim 62, whereby said peptide comprises the amino acid sequence from about amino acid no. 140 to about amino acid no. 337 of SEQ ID No. 1.
- 66. A process according to claim 61, whereby said pathological conditions are any pathological conditions involving the integrin subunit  $\alpha 10$ .
  - 67. A process according to claim 66, whereby said pathological conditions are rheumatoid arthritis, osteoarthrosis or cancer.
- 35 68. A process according to claim 66, whereby said pathological conditions are atherosclerosis or inflammation.

10

35

- 69. A process according to any one of claims 61-68, whereby said cells are chosen from the group comprising chondrocytes, smooth muscle cells, endothelial cells, osteoblasts and fibroblasts.
- 70. A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$ thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.
- 71. A pharmaceutical composition according to claim 70, for use in stimulating, inhibiting or blocking the formation of cartilage, bone or blood vessels.
- 72. A pharmaceutical composition according to claim 15 70, for use in preventing adhesion betweeh tendon/ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue.
- 73. A vaccine comprising as an active ingredient an 20 integrin heterodimer comprising a subunit lpha 10 and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$ , or DNA or RNA coding for said integrin subunit  $\alpha 10$ .
- 74. Use of the integrin subunit  $\alpha 10$  as a marker or 25 target in transplantation of cartilage or chondrocytes.
- 75. A method of using binding entities having the capability of binding specifically to an integrin subunit lpha 10 comprising the amino acid sequence shown in SEQ ID 30 No. 1 or SEQ ID No. 2, or an integrin heterodimer comprising said subunit  $\alpha 10$  and a subunit  $\beta$ , or to homologues or fragments thereof having similar biological activity, for promoting adhesion of chondrocytes and/or osteoblasts to surfaces of implants to stimulate osseointegration.
  - 76. Use of an integrin heterodimer comprising an integrin subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$

10

15

20

25

thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target for anti-adhesive drugs or molecules in tendon, ligament, skeletal muscle or other tissues where adhesion impairs the function of the tissue.

- 77. A method of stimulating, inhibiting or blocking the formation of cartilage or bone, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.
- 78. A method of preventing adhesion between tendon/ ligaments and the surrounding tissue after infection, inflammation and after surgical intervention where adhesion impairs the function of the tissue, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using a integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity, as a target molecule.
- 79. A method of stimulating extracellular matrix synthesis and repair by activation or blockage of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or of the subunit  $\alpha 10$  thereof, or of a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity.
- 80. A method of in vitro detecting the presence of integrin binding entities, comprising interaction of an integrin heterodimer comprising a subunit α10 and a subunit β, or the subunit α10 thereof, or a homologue or fragment of said integrin or subunit, with a sample,
  35 thereby causing said integrin, subunit α10, or homologue or fragment thereof having similar biological activity,

25

30

to modulate the binding to its natural ligand or other integrin binding proteins present in said sample.

- 81. A method of in vitro studying consequences of the interaction of a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, with an integrin binding entity and thereby initiate a cellular reaction.
- 82. A method according to claim 81, whereby the con-10 sequences of said interactions are measured as alterations in cellular functions.
  - 83. A method of using DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof as a target molecule.
- 15 84. A method according to claim 83, whereby a polynucleotide or oligonucleotide hybridises to the DNA or
  RNA encoding an integrin subunit α10 or homologues or
  fragments thereof and whereby said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding
  20 en integrin subunit α1.
  - 85. A method of using a human heterodimer integrin comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit, or a DNA or RNA encoding an integrin subunit  $\alpha 10$  or homologues or fragments thereof, as a marker or target molecule during angiogenesis.
  - 86. A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of stimulating cell surface expression of an integrin heterodimer comprising a subunit  $\alpha 10$  and a subunit  $\beta$ , or the subunit  $\alpha 10$  thereof, or a homologue or fragment of said integrin or subunit  $\alpha 10$  having similar biological activity.





FIGURE 1


2/22

Peptide	Amino acid sequence
1	DNTAQTSAYIQYEPHHSI
2	GPGHWDR
3	AAFDGSGQR
4	FAMGALPD
5	FTASLDEWTTAAR
6	VDASFRPQGXLAP

FIGURE 2

3/22








FIGURE 3




FIGURE 4




FIGURE 5

	CAGGICAGAAACCGATCAGCACTGGAACTCCCCTTCGTCACTCAC	72 -4	CATCCTGCCCAGGGGTGCTGCCTCCATGCCAAGTCCCTCAGCTACTTTTGGCCGAAGTGTGCATGGT R P A Q R I A A A B R P H A L I I V F G R I V D G	1872 195
	GGTETETECTCCCCTTTAACCTGGATCAACATVACCCAGGCCCATTTCCCAGGCCCACCAAGACCCCAATTT	144 19	CONCINCATORISMONACATEMICIATURATURATURATURATURATURATURATURATURATUR	1914 619
	GCATACACTETYTACAACATETTGGGCCGGCCGATGGGGTGGGGTGGGCCCCCCCCCC	216 43	**************************************	2014 6()
	TCAGGGGCGGGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	269 67		2018 667
	CACTINGGIGACIACCACTGGGAATTCATCTCATCCTGCTGGAATAGCACCTGGGGGTGTCTCTGTA R L G O Y O L G H S S N P A V N H N L G N S L L	360 51	COTCOTOGGATCACCANTICTACAGGATCACCGCATGCATGCATGCATGCACTGCAC	2106 691
	CACMAGATOGICATEGGGGATEGATGACTGCTCCCCCCCCCCCCCCCCCCC	432 115	CCATTGATGCCTCGCCACACACTGCCCCTCCCCCCCAATGTCACTGCCAATGTCACTGCAAA	2232 715
	AGTICTECCATATGTCCCCCTGTGGATGCTTCATTCCACCCCACGGAACCCTGCCACCCCACCCCACCCCAACCCCAACCCCAACCCCACCCC	504 139	CARCTACACTTOCATGROCTEGATRACTACACTACCCCCAGTGGCCTTGACGTGACCTTTCCCTTC  0 L B F M V L D T S D Y L B P V A L T V T F A L	2304 739
-	TOCCCANCATACATGGTGTGTGTGTGTGTGTGTGTGTGTGTGAACAACAATGTAGCCCTGGTGTGAAGTTCAG C P T Y H D V V I V I D G S P S L Y P V S E V Q	576 163		2376
	ACCITECTACGGAGGCIGGIAGGGAAACTGIITAITGACCCAGAACAGAIAGAIACAGGIGGGACIGGIACAGIAT T F L B R L V G K L 7 I D P R 0 I 0 V G L V 0 Y	648 . 187	•	2118
	GESCACEDETICACATUAGTELTECCTGGGAGATTTCCCAACGACGAGAGAGTGCTGGAGAGCAGAAGAGGGGGGGAGAGGGGGGAGAGGGGGGAGAGGGGGG	720 211		2520 011
	AACTICATTCOCCGGGGGGGGGGAACAAGACTGCCCAGCAATAATGGTGCCTTCCACAGAGGGTTC R L S R E G R E 7 R 7 R Q A I N V A C 7 E G F	792 235	MAGGAMATGCTTACAMTACGACCTGACTATCATCTTCTCTACAMCCTCCACCTGCCCAGTCTCACCCCCCCCCC	2592
	# ASTENDITUDEATECCECCTCACCCUMCCCTCCCCACCCCACTCCTCCTCTCTCTCATCCCAACTCCCACCC S O S N G G A P S A A R L S V V I D G S S N D	864 259		2444
	COMMONDACTIC COMMON COM	936 283	CCTGTCTTCCMARCHAGGGGACCCAAGGGGACCTTTCCCAAGAGTTTAACTTTAACTCCTCCTCCTCCTCCCCAAGAGTTAACTTTAACTTTAACTCCTCCCCCAAGAGTTAACTTTAACTTTAACTTTAACTCCTCCCCCAAGAGTTAACTTAAACTTAAACTACCAAGAGTAAAAAAAA	2736 883
	GGTCHCTACCTCCGCCGCCAGATCCCAGCTCTTTTCCTCAGAGAAAATTAGAACTATFTCCLAGGATCCA G R Y L R R O N D P 3 S P L R B I R 2 I A 2 D P	1000		2000
	CATCHOCKATTCTTCAMCTCACACATCACCTTCTCAMTCACATTCTCCATCCA	1000	•	2886 931
	ATTITION CONTRACTOR NAME OF THE STATE OF THE	1152 351	CTTCMCCCATAGGGACCCCCCCGGGGGGCGGGGGGGGGGG	2952 955
	TICHTCHTCACTAMAGATCATTTTTTTTTTTTTTTTTTTTTTTTTTT	1224 379		3024
	TOCTICAGGEOCIACCOCTITICCCCCCACEMATGGEACTGGAACAGGTCCCCCCTCCACTGCAG V L C C N R L F P P R N A L E D E F P P A L O	1296 403	CTATURETICEMENTERINETHEMETERACTORINATIONAGECTEACTOMACCOCCA L S L S O V I T W W A S C I V O M L T S P P G P	3096
	MACATICACCTACCTACATATATCTCTTTCTTCATCCTTTTTCCCCCC			3164
		1440	TECHNITION CONTROL AND CONTRACTION CONTROL CONTRACTOR C	3240 1051
	CONTRACTOR AND CONTRACTOR AND	1512 475		3312 1875
	CATCHAMACTERISTICTUACTICCCCCCCCCATGTTCTCCCCCCCCAGACAACGACACACAC	1584		3394
	CITIFATUTE TATESCENCE ACCESSES CONTROL CONTROL CONTROL CONTROL OF THE PROPERTY	1656 523	TOCCTOCATOCICATAGGGAATATACCTGGGAAGGGATTACCTCCCTTCCCTTCCTT	3456 1123
	CHROCKOCCTITECCTTTCCCATCGGGCTCTTCCTGATCTCATCCCAGGTCGTTTTCCTGATCTGGCTCTGG DARFCFANCALLPDLBQDEFADVAV	1720 547		2520 1145
	CONCOCCTICOLALIZACIONACIONACIONACIONACIONACIONACIONACION		4-44-44-4-4-4-4-4-4-4-4-4-4-4-4-4-4-4-	3600
	**************************************	•11	CCTOCATECOCKMONDCCCCTTTTCCTTACCCCCCCCCCCCCCCCCCCCCCCCC	3741 3816 3804

FIGURE 6





FIGURE 7

8/22



B IP: α10 β1

Blot: β1 β1

200 -

97 -

46 -

FIGURE 8





FIGURE 9






FIGURE 10





FIGURE 11


SUBSTITUTE SHEET (RULE 26)

# Human RNA Master blot

Tissue	alo expression	Tissue	a10 expression
Aorta	<del>++++</del>	Thyroid gland	-
Trachea	· +	Salivary gland	-
Lung	++	Spleen	•
Fetal lung	++	Fetal spleen	•
Kidney	++	Thymus	•
Fetal kidney	(+)	Fetal thymus	• .
Heart	(÷)	Peripherial leucocyte	•
Fetal heart	++	Lymph node	-
Spinal cord	++	Appendix	• •
Mammary gland	(+)	Placenta	-
Bone marrow	(+)	Whole brain	•
Small intestine	(+)	Fetal brain	• •
Skeletal muscle	•	Amygdala	-
Liver	•	Caudate nucleus	-
Fetal liver	•	Cerebellum	
Colon	•	Cerebral cortex	-
Bladder	•	Frontal lobe	•
Uterus		Hippocampus	
Prostate	•	Medulla oblongata	-
Stomach	-	Occipitial lobe	•
Testis	-	Putamen	-
Ovary	•	Substantia nigra	-
Pancreas	•	Temporal lobe	-
Piutiatary gland	-	Thalamus	•
Adrenal gland	•	Subthalamic nucleus	-

FIGURE 12









FIGURE 13






FIGURE 14




FIGURE 15a




FIGURE 15b




FIGURE 15c




FIGURE 15d




FIGURE 15e




FIGURE 15f




FIGURE 16




FIGURE 17

International application No.

### PCT/SE 99/00544 A. CLASSIFICATION OF SUBJECT MATTER IPC6: C07K 14/705, A61K 38/17, C07K 16/28 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC6: C07K, A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched SE.DK.FI.NO classes as above Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. Category\* J Biol Chem., Volume 273, August 1998, Lisbet Camper et al, "Isolation, Cloning, and P,X 1-86 Sequence Analysis of the Integrin Subunitalpha 10, a Betal-associated Collagen Binding Integrin Expressed on Chondrocytes, Issue32, page 20383 page 20389 WO 9219647 A1 (THE SCRIPPS RESEARCH INSTITUTE). 1-20,25-29, X 33-44,48-53, 12 November 1992 (12.11.92) 57-62,66-86 21-24,30-32, 45-47,54-56, 63-65 Further documents are listed in the continuation of Box C. See patent family annex. Special categories of cited documents: later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "A" document defining the general state of the art which is not considered to be of particular relevance "E" eriter cocument but published on or after the international filing date "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive "L" document which may throw doubts on priority claim(s) or which is step when the document is taken alone cited to establish the publication date of another citation or other special reason (as specified) document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person stilled in the art document referring to an oral disclosure, use, exhibition or other document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report **30** -07- 1999 14 July 1999 Name and mailing address of the ISA! Authorized officer Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Patrick Andersson/Els

Telephone No. + 46 8 782 25 00

Facsimile No. +46 8 666 02 86

International application No. PCT/SE 99/00544

	LCIVE 33V	70377
C (Continu	ation). DOCUMENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J Cell Biol, Volume 115, No 1, October 1991, Takada Y, Murphy et al, "Molecular cloning and expressin of the cDNA for alpha 3 subunit of human alpha 3 beta 1 (VLA-3), an integrin receptor for fibronectin, laminin, and collagen", page 257 - page 266, Medline abstract acc. no. 92011866	1-20,25-29, 33-44,48-53, 57-62,66-86
A	<del></del>	21-24,30-32, 45-47,54-56, 63-65
<b>X</b>	EMBO J, Volume 8, No 5, May 1989, Takada Y et al, "The primary structure of tha alpha 4 subunit of VLA-4: homology to other integrins and a possible cell-cell adhesion function", page 1361 - page 1368, Medline abstract Acc. no. 89356603	1-20,25-29, 33-44,48-53, 57-62,66-86
A	o-	21-24,30-32, 45-47,54-56, 63-65
X	J.Cell Biiol., Volume 138, No 5, Sept 1997, Lisbet Camper et al, "Integrin alpha2Beta1 Is a Receptor for the Cartilage Matrix Protein Chondroadherin" page 1159 - page 1167	1-20,25-29, 33-44,48-53, 57-62,66-86
A		21-24,30-32, 45-47,54-56, 63-65
X	WO 9425487 A1 (CHILDREN'S MEDICAL CENTER CORPORATION), 10 November 1994 (10.11.94), page 16 - page 21	1-19,28, 33-42,48-52 57-61,66-86
A		20-27,43-47 53-56,62-65

International application No.
PCT/SE 99/00544

ategory *	Citat	ion of document	, with indication, where	appropriate, of the	relevant passages	Relevant to claim No.
A	EP	0330506 A2 30 August	(DANA-FARBER CA 1989 (30.08.89)	NCER INSTITUTI	Ε),	1-86
	-				•	
•						
				·		
•		× .				
				•		
	'					
			•			
						·
		·		•	•	
			•			2
				•		
٠		. •				
					•	
				•		
•						
			• •			
			•			

International application No. PCT/SE99/00544

Box I	Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This inter	rnational search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
ı. 🖂	Claims Nos.: 28-41, 43-69, 74-79, 83-85 because they relate to subject matter not required to be searched by this Authority, namely:
2.	These claims relate to either methods of treatment by therapy or diagnostic methods practised on the human or animal body, see PCT Rule 39.1(iv). Nevertheless, a search has been executed for these claims. The search has been based on the see next page Claims Nos: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This Inte	rnational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark	on Protest The additional search fees were accompanied by the applicant's protest.
	No protest accompanied the payment of additional search fees.

International application No. - PCT/SE99/00544

alleged effects	of the compounds.	
	•	
	•	
		•
·		*
•	•	
	•	
•	•	
	•	·
•		7
*		
·		•
	. •	
	• •	
	•	
	•	
		. 4
		•

Form PCT/ISA/210 (extra sheet) (July1992)

Information on patent family members

01/06/99

International application No.
PCT/SE 99/00544

	itent document in search repor	1	Publication date		Patent family member(s)	Publication date
WO	9219647	A1	12/11/92	AU US US	1896392 A 5310874 A 5589570 A	10/05/94
WO	9425487	A1	10/11/94	AU	6639394 A	21/11/94
EP	0330506	A2	30/08/89	JP US	2003700 A 5583203 A	